

Multi-Scale Nature of the High-Latitude Magnetic Field Fluctuations



Lucia Santarelli¹, Paola De Michelis¹ and Giuseppe Consolini²

¹Istituto Nazionale di Geofisica e Vulcanologia, Roma, Italy- ²Istituto Nazionale di Astrofisica, Roma, Italy




FRAMEWORK

The dynamics of the Earth's magnetosphere in response to changes in solar wind conditions and the interplanetary magnetic field during a magnetic storm and under-storm is the result of both **externally driven processes** and **internal processes**. Recent studies (Alberti et al., 2017) have highlighted the existence of a significant **separation of timescales** in the solar wind-magnetosphere coupling which occurs at approx. **200 minutes** and in particular:


- Long time fluctuations ($T > 200$ min) show a high degree of correlation with the parameters of solar wind and magnetospheric dynamics
- Fluctuations with time-scale $T < 200$ min, although triggered by changes in interplanetary conditions, are mainly dominated by internal processes and are not directly driven (correlated) by the SW/IMF.

DATASET

Data relative to the X and Y component of the Earth's magnetic field recorded at 60 and 20 geomagnetic observatories located in the Northern and Southern Hemisphere, respectively. They are part of the worldwide network of observatories known as INTERMAGNET. We consider a period of some days around the 2015 St. Patrick's magnetic event.



Daily distributions of the X and Y magnetic field components. Maps describe the magnetic field perturbations on the ground due to external sources. The main contributions to this external fields, that produce relevant signatures in magnetic field observations, are the polar ionospheric currents, such as the polar electrojets, and the magnetospheric currents such as the Chapman-Ferraro currents and in particular the ring current.




Method of Analysis: Empirical Mode Decomposition



The EMD method (Huang et al., 1998) is a powerful tool for investigating natural signals that result from non-stationary and non-linear processes. EMD moves from the assumption that most signals are multi-components, involving many scales simultaneously, so that a signal can be considered as a superposition of oscillations from fast to slow ones at a very local level.

Each component (IMF) is defined as any function having:


- 1) the same number of extrema and of zero-crossings in the whole dataset
- 2) a mean value of the envelope defined by the local maxima and the envelope defined by the local minima equals to zero at any point.




$$x(t) = \sum_{i=0}^n IMF_i(t) + res$$



Example of decomposition of the Y component recorded at the CLF observatory. The original signal can be decomposed in 14 different IMFs and a residue. The time scale is increasing with the IMF; each mode has a different mean frequency, which is estimated by considering the mean frequency in the Fourier power spectrum. Taking into account the different values associated with each IMFs, we have decided them in two groups: IMFs with $T < 200$ min and $T > 200$ min.



Analysis and Results



On the left: Daily maps of the ratio between the variance of magnetic fluctuations below 200 minutes and that of the overall signal. The weight of the signal relative to the fast fluctuations shows a dependence on latitude and geomagnetic activity level. At high-latitude the fluctuations on short-time scale have an important weight, in terms of energy, on the total magnetic signal. During the disturbed periods the fast fluctuations have the highest values in the auroral oval. This region is mainly dominated by internal processes and is not directly driven (correlated) by the SW/IMF. A similar behavior characterizes the two hemispheres. However, the ratio between the variance of fast magnetic fluctuations and that of the total signal is higher in the Southern Hemisphere than in the Northern one.

On the right: Daily maps of the ratio between the variance of the slow magnetic fluctuations and that of the overall signal. The weight of the signal related to the fluctuations on a long-timescale (that is triggered by changes in interplanetary conditions) shows a dependence on the latitude and geomagnetic activity level. It is minimum in the auroral regions and maximum moving towards the equator.

Difference between the two hemispheres:.....

Impact on Space Weather: During the disturbed periods the magnetic field fluctuations on short-timescale can play an important role in the total magnetic signal. These fluctuations are mainly dominated by internal processes, although triggered by changes in interplanetary conditions. In order to forecast the magnetic signals it is necessary to develop model that take into account both the changes in interplanetary conditions and in the magnetospheric dynamics.