



OPEN

Excess degassing drives long-term volcanic unrest at Nevado del Ruiz

João Lages^{1✉}, Zoraida Chacón², Julian Ramirez², Alessandro Aiuppa¹, Santiago Arellano³, Marcello Bitetto¹, Julián O. Peña², Diego Coppola⁴, Marco Laiolo⁴, Francesco Massimetti⁴, Lina Castaño², Carlos Laverde², Giancarlo Tamburello⁵, Gaetano Giudice⁶ & Cristian Lopez²

This study combines volcanic gas compositions, SO₂ flux and satellite thermal data collected at Nevado del Ruiz between 2018 and 2021. We find the Nevado del Ruiz plume to have exhibited relatively steady, high CO₂ compositions (avg. CO₂/S_T ratios of 5.4 ± 1.9) throughout. Our degassing models support that the CO₂/S_T ratio variability derives from volatile exsolution from andesitic magma stored in the 1–4 km depth range. Separate ascent of CO₂-rich gas bubbles through shallow (< 1 km depth), viscous, conduit resident magma causes the observed excess degassing. We infer that degassing of ~ 974 mm³ of shallow (1–4 km) stored magma has sourced the elevated SO₂ degassing recorded during 2018–2021 (average flux ~ 1548 t/d). Of this, only < 1 mm³ of magma have been erupted through dome extrusion, highlighting a large imbalance between erupted and degassed magma. Escalating deep CO₂ gas flushing, combined with the disruption of passive degassing, through sudden accumulation and pressurization of bubbles due to lithostatic pressure, may accelerate volcanic unrest and eventually lead to a major eruption.

Volcano monitoring and eruption forecasting have greatly benefited from recent technological advances that allow high temporal resolution measurements of volcanic gas compositions and fluxes. Volcanic gases measured at the surface are the only direct chemical probe of magma at depth and may, by their composition and/or flux, indicate movement of magma toward the surface, changes in the permeability of the shallow conduit system, or pressurization of the magma column beneath a lava dome^{1–4}.

Therefore, improving geochemical monitoring infrastructures, and enabling real-time analysis and interpretation protocols, are paramount to our understanding of pre- and syn-eruptive behavior of persistently degassing volcanoes and to mitigate the risk they pose to vulnerable communities.

Nevado del Ruiz, in Colombia, is a 5.321 m-high glacier-clad andesitic volcano in the Cordillera Central of the northern Andes. The volcano erupted numerous times during the Holocene. Its 20th century eruptive history was marked by a period of unrest beginning in late November 1984 with a sharp increase in fumarolic activity⁵. It culminated with an eruption on 13 November 1985, which generated large lahars and killed more than 23,000 people^{5–7}.

More recently, deformation was noted in 2007, while seismicity and SO₂ emission rates started increasing in 2010, with SO₂ fluxes associated with small eruptions in May and June of that year^{8–10} reaching levels in excess of 20,000 t/d in 2012. In the meantime degassing rates between 2005 and 2015 remained high at Nevado del Ruiz, with satellite data showing an average SO₂ flux of ~ 1,074 t/d¹¹ leading up to elevated deformation and two peaks in lava dome extrusion rate: a first short-lived pulse in November 2015 and a second lasting most of 2016. By the beginning of this study, extrusion rates had decreased to 0.02 m³/s (February 2018). These continued to decline until February 2019, when the dome forming eruption eventually ended^{12,13}.

Lava domes are structures that result from the extrusion and accumulation of extremely viscous, quasi solid, lava that are commonly formed at andesitic stratovolcanoes like Nevado del Ruiz. Explosive eruptions at lava domes are thought to be caused by spatial and temporal changes in their permeability and of their ability to exsolve and release volatiles^{14,15}. Volcanic gas observations, especially if combined with thermal satellite observations¹⁶, are thus especially relevant to understanding lava dome activity and behaviour^{17,18}. For Nevado del Ruiz, no information on the fluxes of other major volatile species, such as H₂O and CO₂, was available until 2017, when the first discontinuous measurements started¹⁹.

¹Dipartimento DiSTeM, Università Degli Studi Di Palermo, Palermo, Italy. ²Servicio Geológico Colombiano, Observatorio Vulcanológico y Sismológico de Manizales, Manizales, Colombia. ³Department of Space, Earth and Environment, Chalmers University of Technology, Gothenburg, Sweden. ⁴Dipartimento Di Scienze Della Terra, Università Di Torino, Torino, Italy. ⁵Istituto Nazionale Di Geofisica E Vulcanologia, Sezione Di Bologna, Bologna, Italy. ⁶Istituto Nazionale Di Geofisica E Vulcanologia, Osservatorio Etneo, Catania, Italy. ✉email: joaopedro.nogueiralages@unipa.it

This study reports systematic volcanic gas observations (CO_2/SO_2 ratios, CO_2 and SO_2 fluxes) taken in 2018–2021, a period of declining dome extrusion rates and negligible deformation. Nonetheless, seismicity, gas and ash emissions remained prevalent throughout this study¹³. Our aim is to present a model of the processes sustaining the persistent degassing, and to identify the mechanisms through which volcanic activity may escalate during periods of prolonged (slow) unrest²⁰.

Results

Volcanic gas compositions

Our results are based on volcanic gas records streamed by a fully autonomous MultiGAS^{21,22} station. The instrument was deployed at Nevado del Ruiz between 2018 and 2021, on the northwest flank of the volcano at an altitude of 4832 m a.s.l. (4.90°N , -75.34°W ; Fig. 1). The data yield¹⁹ average CO_2/SO_2 ratios of 5.4 ± 1.9 (2.8–14.3, $n = 220$; Fig. 2A; see “Methods”). H_2S concentrations were rarely detected at > 1 ppm levels, and the $\text{H}_2\text{S}/\text{SO}_2$ ratios are typically < 0.1 . Volcanic H_2O signal (above atmospheric background; see “Methods”) is resolved in only 25 acquisitions, due to the very high background (ambient) air H_2O concentrations (up to 16,000 ppm) recorded at such altitudes. These yield $\text{H}_2\text{O}/\text{SO}_2$ and $\text{H}_2\text{O}/\text{CO}_2$ ratios averaging at 32.8 (range, 9.1–56.7) and 3.9 (2.6–6.5),




Figure 1. Satellite image of Nevado del Ruiz showing the location of MultiGAS ($n = 1$) and NOVAC stations ($n = 5$) used in this study. **(A)** Distribution of SO_2 Max concentrations recorded by the MultiGAS station between 2018 and 2021; and **(B)** Wind direction data from the NOVAC network, showing good agreement between the location of the permanent MultiGAS station and the predominant wind direction. On the right, photos taken from the monitoring webcams between 2018 and 2021 are courtesy of the *Observatorio Vulcanológico y Sismológico de Manizales (Servicio Geológico Colombiano)*.




Figure 2. (A) CO_2/SO_2 compositions (molar); and (B) Daily SO_2 fluxes (averages in t/d; NOVAC Network). (C) CO_2 fluxes are derived from the combination of SO_2 flux estimates and MultiGAS measurements (see “Methods”). The red lines represent a 10-pt. average (A–C).

respectively. From these measurements, we estimate the average composition of the plume at 84.7 mol% H_2O , 12.0 mol% CO_2 , 2.8 mol% SO_2 , 0.1 mol% H_2S , and 0.4 mol% H_2 . Uncertainties in gas ratios measured by the Multigas are reported in Supplementary Table 1 and are far lower than the variations reported in our time series.

SO_2 fluxes

Daily average SO_2 fluxes (see “Methods” for data selection criteria and details on daily statistics of SO_2 emission rates), obtained by the local NOVAC²³ network of 5 scanning spectrometers between 2018 and 2021, oscillated between 58 and 4617 tons/day, with an average of 1568 tons/day (Fig. 3A). This confirms the sustained degassing activity of Nevado del Ruiz during the investigated time interval. Annual averages show small variations, especially between 2018 (~1457 tons/day) and 2019 (~1590 tons/day). Four out of the 5 stations yield somewhat similar yearly averages, ranging from ~2910 (Bruma) to ~4031 t/day (Azufrado/Olleta), thus attesting for the uninterrupted degassing and somewhat unvarying activity at the Arenas crater.

Volcanic radiative power

In the temporal interval investigated, the MIROVA²⁴ system detected intermittent thermal anomalies, with a Volcanic Radiative Power (VRP) baseline below 5 MW (Fig. 4A). These relatively low VRP values attest for the overall mild lava extrusion activity registered at Nevado del Ruiz between 2018 and 2021, coupled with continuous high-temperature degassing. Periods of dome extrusion (e.g., Jan–Apr 2020) are clearly detected by MIROVA as VRP maximum values of up to 16.7 MW (see supplementary Table 1–2 for detailed thermal outputs).




Figure 3. (A) Relative frequency distribution of SO_2 fluxes between 2018–2021 (NOVAC Network; time series shown in Fig. 2B); (B) The same data is shown for days in which ash emissions were detected (total of events/days = 51; see “Methods”). Note that in the occurrence of ash emissions approximately 59% of SO_2 fluxes fall below the 3-year SO_2 flux average of ~ 1570 t/d.

Discussion

Shallow versus deep magmatic gas signature

The near absence of H_2S in the gas plume (avg. ~ 0.1 mol%) suggests negligible hydrothermal contributions to volcanic gas compositions measured at Nevado del Ruiz between 2018 and 2021. The magmatic nature of the measured gas is additionally supported by the relatively low H_2O concentrations (maximum 92 mol%). Therefore, we focus on the temporal variations of plume CO_2/SO_2 ratios (Fig. 2A), and on the fluctuations of SO_2 and CO_2 fluxes (Fig. 2B, C). The in-plume abundances of CO_2 and SO_2 both exhibit significant temporal variations. The relatively high CO_2/SO_2 ratio range (5.4 ± 1.9) confirms the C-rich nature of Nevado del Ruiz magmatic fluids, interpreted^{19,25} as originating from the recycling of subducted carbonate-rich sediments in the region²⁶ (see Aiuppa et al., 2017 for detailed assessment of the relationship between along-arc CO_2/SO_2 ratios and subduction sediment compositions). Above average CO_2/SO_2 ratios are unlikely to be caused by the scrubbing of volcanic SO_2 (a process that can cause CO_2/SO_2 ratios to exceed typical magmatic values²⁷) for two main reasons. Firstly, a typical driver of magmatic S scrubbing is the interaction of deeply ascending magmatic fluids with hydrothermal fluids/ground-water, whereby the conversion of SO_2 to H_2S should occur; this is not observed at Nevado del Ruiz, given the negligible amounts of H_2S measured. Secondly, at andesitic dome-forming volcanoes, SO_2 scrubbing should be favored in phases when cooling and/or mineral deposition in fractures and pores in the dome carapace²⁸ prevail. If this was the case, then high CO_2/SO_2 ratios should systematically be associated with reduced SO_2 fluxes (reduced SO_2 fluxes have been detected prior to explosion at some dome-forming




Figure 4. (A) Volcanic Radiative Power (VRP; in MW) retrieved from MODIS (blue markers), and associated cumulative thermal energy (Volcanic Radiant Energy; VRE in Joule). High VRP measurements (Jan–Apr 2020) are highlighted by the shaded red area, and also on the inset for comparison with extrusive events of 2015–2016. (B) 2018–2021 Extrusion rates reported in Ordoñez et al. (ref¹³ for details). On the inset of B, note the good agreement between VRP data (2012–2021; this study) and extrusion rates, especially for the two extrusion rate peaks detected in November 2015 and for most of 2016¹³. (C) Cumulative volumes of degassed (in Mm^3), thermally radiant (as V_{Thermal}) and extruded magma¹³ (see “Methods” for details on calculations).

volcanoes, interpreted as caused by the decreasing in permeability of the main degassing pathways¹⁷). However, at Nevado del Ruiz, we observe persistently high SO_2 fluxes (Fig. 2B) that attest to an overall permeable dome, allowing efficient escape of magmatic gases to the atmosphere. We also find no significant correlation between the timing of the summit ash explosions and SO_2 fluxes (Fig. 3). If we concentrate on the days in which at least one explosion is observed (Fig. 3B), we note that in only 59% of these the daily recorded fluxes are below the 2018–2021 average (41% of the days with explosions recorded higher-than-average SO_2 fluxes). We caution that we are here interested in long-term (daily to yearly) degassing trends rather than in the driving mechanisms of ash explosions, and we cannot exclude short-term (minutes to tens of minutes) drops in SO_2 emissivity occur prior to individual explosions (as observed elsewhere¹⁷) that are not resolvable at the scale of our observations here. In our context, we conclude that clusters of explosions can occur in periods of either reduced (125–1000 t/d) or augmented (2000–3000 t/d, and up to 4617 t/d) daily SO_2 emission rates. Ultimately, we see no obvious link between compositional changes and shallow processes (scrubbing, dome permeability drop), and we find more likely that the temporally changing CO_2/SO_2 ratios are linked to magmatic processes, and potentially to a variable input of deeply rising CO_2 -rich fluids²⁹ into the shallow magma plumbing system feeding the dome.

Modelling magmatic degassing requires an understanding of volatile contents in the Nevado de Ruiz parental melts. Stix et al., (ref.³⁰) analysed juvenile material erupted at Nevado del Ruiz in November 1985 and September 1989. The authors argued that the wide range of SiO₂ contents (62.4–76.6 wt%) observed in melt inclusions implies two distinct magmas are at play, one more evolved than the other. This hypothesis is frequently invoked in the post-1985-eruption literature^{31–33}. Here we interpret our volcanic gas compositions by using, as proxy for the parental (undegassed) melt composition, the measured volatile contents (2.45 wt% H₂O and 440 ppm S) in the less evolved (62.4 wt% SiO₂) melt inclusions³⁰. The CO₂ parental melt concentration has not been characterized at Nevado del Ruiz using melt inclusions. We hence consider a range assumed to be characteristic of initial CO₂ contents in arc magmas by Plank & Manning, 2019 (1200 ppm, ref.³⁴) and Wallace, 2005 (3000 ppm, ref.³⁵).

With these initial input parameters, we use a volatile saturation code^{36,37} to calculate the pressure-dependent evolution of the magmatic gas phase exsolved from Nevado del Ruiz magmas upon their ascent and decompression (Figs. 5 and 6). Our simulations are performed in both closed- and open-system conditions (250–0.1 MPa range) at a constant temperature of 900 °C (1173 K)³², and exploring a redox range of 0.5 log units below the nickel-nickel oxide (NNO) buffer (see supplementary Table S4 for detailed input parameters). Note that the large mismatch between degassing and erupted magma volumes (see below) requires gases are separated from melt (e.g., that open system prevails) at some point in the magma ascent/decompression path. However, as the depth/pressure of closed-to-open degassing transition is undetermined, we examine the full closed and full open conditions separately as two end-member scenarios.

Results (Fig. 5) show that the modeled open- and closed-system degassing trends match well the range of gas (this study, Fig. 5B, D) and melt compositions³⁰ (Fig. 5A, C) observed at Nevado del Ruiz. We can therefore infer the pressures/depths of gas–melt separation (final equilibration) in the plumbing system by comparing the modeled and observed gas compositions (Fig. 6).

Under closed-system conditions the melt becomes volatile saturated at approximately 250 MPa and our lower/upper range of volcanic gas CO₂/S_T ratios would imply equilibration pressures of approximately 30–100 MPa (~1–4 km; Fig. 6). Beyond ~30 MPa pressures the magmatic gas phase would evolve to CO₂/S_T compositions lower than those measured in the gas plume (Fig. 6). On the other hand, for the open-system scenario, CO₂/S_T derived pressures/depths range from ~20 to 93 MPa (~0.8 to 3.7 km).

Our gas-inferred depth range corresponds to those inferred from melt inclusion entrapment conditions³⁰, and to the seismically identified active magma volume¹⁰. Combined with existing knowledge on the shallow Nevado del Ruiz plumbing system^{10,30}, our results identify a main magma storage region in the 1–4 km range, where ponding magma crystallizes (eventually evolving from andesite to dacite), and where gas–melt separation takes place that sustain magmatic gas emissions at the surface. Here, the upper range of our volcanic gas compositions (CO₂/S_T upper range 5.4–7.3; S_T stands for total sulfur, and corresponds to the sum SO₂(g) and H₂S(g)) may correspond to the roots of such magma storage zone (90–100 MPa pressure; Fig. 6), where separate ascent of deeply-derived CO₂-rich gas (CO₂-flushing) starts, eventually followed by separate gas bubble ascent and/or further bubble re-equilibration (1–3 km-depth range). In this interpretation, the shallowest (<20–40 MPa) portion of the plumbing system would then be occupied by relatively stationary (or poorly mobile), viscous andesitic magma, a very small fraction of which is finally extruded as a dome. In this portion of the reservoir, below-average volcanic gas compositions derive from low-pressure re-equilibration and partial CO₂ loss from the melt.

Therefore, we argue that the intermittent resupply of the shallow resident conduit magma with more volatile-rich magma (rising from deep) does play a crucial role in sustaining the long-lasting degassing activity of the magmatic column (in addition to causing the brief excursions of gas compositions toward higher CO₂/S_T compositions). In addition, at low confining pressures and high magma viscosities, there may be sufficient strain at the conduit walls to induce brittle failure, with gas loss along permeable channels³⁸ (Fig. 6). Such lines of evidence corroborate a multistage model of magma transport and degassing, with alternating periods of magma ascent and ponding³⁰.

Dynamics of shallow ponding conduit magma

Assessments of magma balances (e.g., degassed versus extruded) can provide further constraints on magma feeding processes into the shallow Nevado del Ruiz magmatic system. The volume of degassed magma between 2018 and 2021, inferred from the measured SO₂ fluxes and knowledge of parental melt S content (see “Methods”), is ~974 mm³ (Fig. 4C). Additionally, we estimate a mean MIROVA-derived extrusion rate (TADR; see “Methods”) of 0.37 m³/s (andesite), which is considerably higher than that (0.02 m³/s) reported by Ordoñez et al. (ref.¹³) for the 2018–2021 period (Fig. 6). Following the equations provided in Coppola et al., 2013 (ref.³⁹; see also “Methods”), we calculate that the thermal output recorded requires surface emplacement (extrusion) of about 27.5 mm³ of magma (V_{thermal}; Fig. 4C), which is approximately 50 times higher than that of the volume extruded (0.56 mm³)¹³ during that period.

In other dome-forming volcanoes (e.g. Sabancaya¹⁶ and Popocatepetl¹⁸), V_{Thermal} > V_{Extruded} unbalances have been ascribed to an “excess radiation” process whereby the majority of the thermal anomalies (reported as VRP) were sourced by additional processes other than surface dome extrusion^{16,18}. We caution that, at Nevado del Ruiz, the latter may be somewhat underestimated, considering the cycles of dome building and partial destruction (potentially sudden) can be relatively short, and hence difficult to capture with the relatively low temporal resolution measurements reported by Ordoñez et al. Short-lived dome (emplacement/destruction) cycles may, in fact, explain (i) the relatively mild explosive activity of the arenas crater and the lack of a major explosive event since the beginning of the long-lasting unrest; and (ii) the relatively efficient (partial) clearing of the top of the magma column allowing for the conduit to sustain a high level of gas permeability.

In any case, the large unbalance between magma input (10 m³/s) and output (extrusion, 0.02 m³/s) rates, shown in Fig. 4C and schematically illustrated in Fig. 6, indicates that only about 0.2% of the intruded magma




Figure 5. On the left, H₂O (wt.%) vs S (ppm) in melt inclusions from 1985–1989 eruptive products. Lines illustrate the model-predicted³⁶ dissolved H₂O and S contents in the melt along the modelled (A) open- (in blue) and (C) closed-system (in red) degassing paths in the 250–0.1 pressure range (see supplementary Table 4 for full input parameters). On the right, triangular plot comparing model-predicted (lines) and measured gas compositions in the H₂O/10-CO₂*5-S_T*10 magmatic system for both open- (B) and closed-system (D) degassing. Note that model runs fit at large both melt inclusion data³⁰ and measured gas compositions.

finally reaches the surface. Unbalance between supplied and erupted magma (and the notions of excess degassing and thermal radiation highlighted in our dataset) is typical of open-vent-like-behavior and may indicate that, throughout this study, activity (slow unrest) at Nevado del Ruiz was driven by degassing of unerupted magma (see also ref.¹⁶).

We have so far established that the existing lava dome at Nevado del Ruiz is connected to deeper reservoirs (e.g., 1–3 km depth³⁰) through a gas-permeable volcanic conduit (e.g., ref.⁴¹). On the other hand, magma supply rate and erupted magma volume suggest that less 1% of the intruded magma reaches the surface (see above). If such significant volumes of degassed magma were to be stored at shallow depths beneath Nevado del Ruiz (i.e., in the upper 2 km), measurable deformation was to be expected. On the contrary, the local *Observatorio*




Figure 6. Schematic model of shallow conduit processes in play at Nevado del Ruiz, highlighting the discrepancy between magma input (this study) vs output¹³ rates for the 2018–2021 period. Model-predicted, pressure-dependent evolution of the CO_2/S_T ratio in the magmatic gas coexisting with a Nevado del Ruiz-like melts is shown for the model runs in Fig. 5. Note that the exsolution depths yielded by our model runs agree with reservoir depths inferred in the literature^{10,38}.

Vulcanológico y Sismológico de Manizales reported no significant anomalies (not to the scale of the volumes of non-erupted magma) between 2018 and 2021.

We, therefore, argue against the possibility that large volumes of magma are being stored at shallow levels within the edifice. Models of convecting magma columns⁴⁰ have been evoked to explain excess degassing and thermal radiation associated with dome-forming activity at andesitic volcanoes^{16,18,42}. At Nevado del Ruiz, due to significant degassing-induced crystallization in the shallow part of the conduit (Fig. 6), bimodal flow and magma convection may not occur as efficiently as in low-viscosity mafic systems, especially as magma becomes more evolved and stagnant at shallower levels. During the early stages magma crystallization and bubble formation, some extent of counterflows of ascending (non-degassed) and descending (degassed) magma may coexist in the deep (> 3 km) volcanic conduit, therefore boosting the continuous supply and recycling of deep magmatic fluids between reservoirs (Fig. 6).

In the shallower regions of the conduit, gas–melt separation is likely driven by cooling and crystallization of stagnant, viscous andesitic magma. This process concentrates volatiles in the remaining melt phase and eventually causes them to exsolve into bubbles, which in turn propels the steady degassing behavior and gas compositions observed between 2018 and 2021, and permit large fractions of reservoir volatiles to be released without major eruption. Deeper reservoirs connected to shallower regions by dykes provide occasional inputs of CO_2 -rich magma (CO_2 -flushing) which may disturb normal rates of magma ascent and degassing and cause conduit overflow, resulting in the extrusion events recorded in this study.

The eruptive cycle of Nevado del Ruiz volcano: clues on the possible activity escalation of a slow and steady system

Periods of enhanced activity, such as higher rates of dome growth or explosive activity, are common at volcanoes such as Nevado del Ruiz. However, our results corroborate that “slow” silicic systems can eventually maintain a steady-state volcanic activity behavior for years, without ever transitioning into a climatic phase²⁰. Between 2018 and 2021, this “steady-state” behavior has resulted from a complex but overall “balanced” interplay between inputs of volatile-rich magma, shallow magma crystallization and degassing, and dome extrusion, which has only produced relatively mild explosive activity. Similar slow-unrest systems²⁰, of equally evolved magma compositions, such as Popocatepetl^{17,18}, in Mexico, and Sabancaya¹⁶, in Peru show similar longevity in their unrest periods and surface activity. Therefore, a crucial question for these systems, and in particular of Nevado del Ruiz, is how, and over what timescales, volcanic activity can escalate into more voluminous/energetic eruptive events of potential threat to vulnerable communities.

The months preceding Nevado del Ruiz’s catastrophic November 13, 1985 eruption were characterized by minor ash emission events that culminated in a relatively small eruption (Volcanic Explosivity Index, VEI = 3)^{5,43}. Juvenile scoria and pumices were erupted³¹ and about 90 kt of SO₂ released⁴⁴, suggesting that the eruption was in fact magmatic and not phreatic⁴⁵. Giggenbach et al. (ref.⁴⁶; see also ref.⁴⁷) reported on an extensive hydrothermal system beneath the volcano, which is manifested today entirely through springs and fumaroles spread throughout the large periphery of the volcano. Our volcanic gas data, however, shows that the present high gas and heat fluxes have most likely boiled off any meteoric water and potentially decoupled the hydrothermal and magmatic systems of Nevado del Ruiz. If Nevado del Ruiz is to sustain its current levels of unrest, the origin and nature of a future major eruptive event is therefore likely to be magmatic.

Given the catastrophic consequences of the November 1985 eruption⁶, we must attempt to correlate the pre-, syn- and post-eruptive observations of the historical event with the current unrest signals. We emphasize two major findings: (i) Banks et al. (ref.⁴⁸) reported no deformation and therefore lack of significant intrusive activity prior to and during the 1985 eruptive period; and (ii) the amount of “new” magmatic material produced during the November 1985 eruption was disproportionately small to account for the large amounts of SO₂⁴⁹ released then and over subsequent periods (see ref.⁴³). Based on our findings, a large degassing excess and a lack of deformation are distinctive features of present-day activity, although no major eruption has yet occurred. Our conceptual model (Fig. 6) accounts for different evolving magmas³⁰ at shallow depths, which degas extensively over time. The same magma regions were likely involved as source of the large amounts of pre- and syn-eruptive passive degassing observed from 1985 to 1990 and beyond⁴³.

The mechanisms of gas/magma transfer within the shallow magma plumbing system, and between the shallow and deep magmatic systems, are difficult to constrain. However, our results suggest that crystallization-induced (evolved magma) and CO₂-rich gases (from deep) are necessary to explain the range of CO₂/SO₂ compositions measured at the surface. Depending on magma physical properties (e.g., viscosity, vesicularity, and percentage of interconnected vesicles), each of them can dominate at specific depths or time¹⁷. In the current degassing unrest, in particular, phases of enhanced CO₂ flushing can be detected as periods of escalating CO₂ surface release (blue shaded areas; Fig. 7). Increased gas flushing may render the shallow ponding magma more buoyant, eventually leading to occasional events of dome extrusion (red shaded areas, as identified from increasing magma output/input ratios; Fig. 7) once the top of the magma column overflows. Carbon dioxide (CO₂) flushing, in particular, may play a crucial role in governing degassing behavior over time. While at present shallow ponding magma may be sourcing the enhanced degassing rates recorded at Nevado del Ruiz (green shaded areas; Fig. 7), ascent of voluminous CO₂-rich deep gas amounts in the conduit may eventually cause eruption⁵⁰. Volcanic gas release through permeable conduit walls and dome during times of passive degassing may be disrupted by sudden




Figure 7. Magma output/input ratio (2018–2021), and CO₂ and SO₂ cumulative masses distinguish periods dominated by CO₂ flushing (deep) and steady-state degassing, with occasional overflow (minor dome extrusion events) of the magma column. Note that, given the good agreement between extrusion rates¹³ and VRP data between 2012 and 2021 (see Fig. 4B), we use here magma output rate as TADR (in m/s; see “Methods”) to identify periods of higher extrusion rates between 2018 and 2021.

accumulation and pressurization of bubbles due to lithostatic pressure that tends to compact and close the system¹⁷. The combination of both processes may culminate in a major eruption.

Therefore, monitoring the composition and mass flux of volcanic gases is critical for fully informed forecasting efforts. However, the challenges of real-time measurements of volcanic gas compositions at volcanoes such as Nevado del Ruiz are exacerbated by extreme low ambient temperature conditions and high level of volcanic activity. Nonetheless, our study attests to the advantages of combining composition, flux and satellite remote sensing measurements to efficiently address the dynamics of shallow magma transfer and extrusion at strongly degassing volcanoes. Moreover, by monitoring the Nevado del Ruiz volcanic degassing behavior over the 3-year period, this study crucially distinguishes several activity phases (e.g., CO₂ gas flushing, dome extrusion, persistent open-conduit degassing) within the recent unrest cycle of Nevado del Ruiz, while highlighting their specific chemical and thermal patterns to future risk assessment efforts.

Methods

Permanent MultiGAS station

During operation, the MultiGAS^{21,22} measured in-plume concentrations of CO₂, SO₂ and H₂S at 1 Hz. The permanent station worked for 4 30-min cycles every day between 2018 and 2021, at 0:00, 6:00, 12:00, 18:00 (UTC time). For details on calibration and sensor range see ref.⁵¹. Ambient pressure, temperature and relative humidity were also measured, which allowed calculation of in-plume H₂O concentrations using the Arden Buck equation⁵² (Supplementary Table 3). CO₂/SO₂ and H₂O/SO₂ ratios (supplementary Tables 1 and 3) correspond to the slope of a best-fit regression line of the concentrations (in ppm) of both species in the selected temporal window (Ratiocalc⁵³). Results (Fig. 2) are only reported for temporal windows in which the SO₂ concentration was above the 5 ppmv threshold, and in which correlations between CO₂ and SO₂ and H₂O and SO₂ exceed an R² of 0.6. Despite the daily measurement routines, our volcanic gas dataset is limited to days in which wind direction favored the southwest sector of the volcano, where the sector Bruma is located (see Fig. 1). For instance, between 2018 and 2019, 1725 acquisitions (30-min each) were successfully transferred via telemetry from Bruma to OVSM, and subsequently processed at the University of Palermo. Approximately 67% of these acquisitions registered SO₂ concentrations above instrument noise (>0.2 ppmv), but only about 23% recorded SO₂ levels ≥ 5 ppm (the minimum concentration threshold here considered above which the plume is sufficiently “dense” to allow for compositional and CO₂ flux estimates; Fig. 2). Error are expressed as the standard error of the regression analysis and subsequent error propagation, error on inferred flux propagate error on the SO₂ fluxes and gas ratios.

Daily SO₂ flux estimates

Sulfur dioxide emissions from Nevado del Ruiz are measured daily by scanning UV spectrometer systems installed through the Network for the Observation of Volcanic and Atmospheric Change project^{23,54}. This network includes 5 different scanning locations, Bruma (4.90; -75.33, 4878 m a.s.l.), Alfombrales (4.88; -75.35, 4458 m a.s.l.), Azufrado/Olleta (4.89; -75.35, 4909 m a.s.l.), Inderena/El Camion (4.96; -75.37, 4016 m a.s.l.) and Recio 3 (4.86; -75.33, 4665 m a.s.l.; see map of Fig. 1) that provide plume scans at virtually all wind directions. The NOVAC scanning mini-DOAS (differential optical absorption spectroscopy; see ref.⁵⁵) instruments scan the sky continuously during daylight hours to measure the integrated absorption of UV light by SO₂ in the plume. These are then combined with meteorological information to derive daily statistics of total SO₂ emissions. Wind speed and direction are acquired from local meteorological models from IDEAM (<http://www.ideam.gov.co/>). Daily SO₂ flux estimates are combined here with in tandem CO₂/SO₂ gas ratios (converted from molar ratios to mass based on concentration ratios) measured by the permanent MultiGAS station to derive CO₂ flux budgets between 2018 and 2021. The SO₂ flux dataset assembled over the years by the OVSM highlights a dependence on wind patterns. Specifically, between May and October, westwards plume directions allow ideal scanning geometries for 4 out of the 5 stations (located on the west flank of the volcano). This ultimately translates into higher estimated fluxes comparing to periods during which the plume may become undetected in more than one scan (due to unfavorable transport directions). We here consider only SO₂ flux measurement scans with complete coverage of the plume (completeness > 0.8), in order to minimize the effect of wind direction in our daily SO₂ flux estimates.

CO₂ fluxes

We derive daily averaged CO₂ fluxes (in t/d; Fig. 2C) by combining CO₂/SO₂ ratios (MultiGAS station) and SO₂ fluxes (NOVAC network), as: CO₂ flux = CO₂/O₂ × SO₂ flux (t d⁻¹).

Sulfur flux (in kg/s) is calculated from the following:

$$S_{\text{flux}} (\text{kg s}^{-1}) = \frac{M(\text{S})}{M(\text{SO}_2)} \times \left(\frac{\text{SO}_2 \text{ flux (intd}^{-1}) \times 1000}{24 \times 60 \times 60} \right).$$

Volcanic radiative power (MODIS)

MIROVA²⁴ (Middle InfraRed Observation of Volcanic Activity; www.mirovaweb.it) algorithm allows to detect, locate and quantify volcanic hotspots, measuring the heat flux radiated by hot (> 300 °C) volcanic features (VRP ± 30%, Fig. 4A, inset of Fig. 4B). This approach provides the VRP time series (and its associated Volcanic Radiative Energy, VRE; Fig. 4A) recorded at Nevado del Ruiz between 2018 and 2021 and prior. **Volumes of radiating magma (V thermal; Fig. 4C)** are retrieved from the thermal approach, and are related to the measured radiant energy (VRE)³⁹ through:

$V_{\text{Thermal}} = \frac{VRE}{C_{\text{rad}}}$, where C_{rad} is an empirical coefficient that takes into account the effective rheology of the emplacing lava body³⁹. For Nevado del Ruiz we used a silica content of 62.4 wt%³⁰, obtaining a C_{rad} of 2.1×10^6 J/m³.

The **volume of degassed magma (V degassed;** Fig. 4C) takes into account the measured S flux (see above) and calculations of magma input rates^{40,56}. It is obtained from the following equation:

$$V_{degassed} = \frac{S_{flux}(in\ kg\ s^{-1})}{\Delta XS \times \rho_m \times \phi},$$

where ΔXS is the S volatile loss, derived from the difference between parental melt volatile content (440 ppm in melt inclusion data³⁰) and the residual S content in the groundmass (as low as 70 ppm³⁰); ϕ is the crystal fraction, assumed for Nevado del Ruiz magmas to be around 40%; and ρ_m is the melt density of the magma (2227 kg/m³⁽³⁰⁾).

Same estimates of **magma input rates** are used in Fig. 7. Instead, given the good agreement between VRP and output rates shown in Fig. 4B (and inset), output rates in Fig. 7 are calculated as time-averaged lava discharge rates (TADR)³⁹, by considering the following: $TADR = \frac{VRP}{C_{rad}}$.

Ash events record

Ash emission events were registered by the *Observatorio Vulcanológico y Sismológico de Manizales (Servicio Geológico Colombiano)* through observation of webcam video recordings and reports from local communities.

Received: 8 August 2023; Accepted: 4 January 2024

Published online: 12 January 2024

References

1. Aiuppa, A. Volcanic-gas monitoring. In *Volcanism and Global Environmental Change* 81–96 (Cambridge University Press, 2015).
2. Aiuppa, A. *et al.* A CO₂-gas precursor to the March 2015 Villarrica volcano eruption. *Geochem. Geophys. Geosyst.* **18**, 2120–2132 (2017).
3. Edmonds, M., Herd, R. A., Galle, B. & Oppenheimer, C. M. Automated, high time resolution measurements of SO₂ flux at Soufrière Hills Volcano. *Montserrat. Bull. Volcanol.* **65**, 578–586 (2003).
4. Edmonds, M. Geochemical monitoring of volcanoes and the mitigation of volcanic gas hazards. *Forecast. Plan. Volcan. Hazards Risks Disasters* **2**, 117–151 (2021).
5. Hall, M. L. Chronology of the principal scientific and governmental actions leading up to the November 13, 1985 eruption of Nevado del Ruiz, Colombia. *J. Volcanol. Geoth. Res.* **42**, 101–115 (1990).
6. Voight, B. The 1985 Nevado del Ruiz volcano catastrophe: anatomy and retrospection. *J. Volcanol. Geoth. Res.* **42**, 151–188 (1990).
7. Thouret, J. C., Cantagrel, J. M., Salinas, R. & Murcia, A. Quaternary eruptive history of Nevado del Ruiz (Colombia). *J. Volcanol. Geoth. Res.* **41**, 225–251 (1990).
8. Castaño, L. M. *et al.* Continuous monitoring of the 2015–2018 Nevado del Ruiz activity, Colombia, using satellite infrared images and local infrasound records. *Earth Planets Space* **72**, 81 (2020).
9. Federico, C. *et al.* Vapour discharges on Nevado del Ruiz during the recent activity: Clues on the composition of the deep hydrothermal system and its effects on thermal springs. *J. Volcanol. Geoth. Res.* **346**, 40–53 (2017).
10. Vargas, C. A. *et al.* Breathing of the Nevado del Ruiz volcano reservoir, Colombia, inferred from repeated seismic tomography. *Sci. Rep.* **7**, 8–13 (2017).
11. Carn, S. A., Fioletov, V. E., Mclinden, C. A., Li, C. & Krotkov, N. A. A decade of global volcanic SO₂ emissions measured from space. *Sci. Rep.* **7**, 1–12 (2017).
12. Londono, J. M. & Galvis, B. Seismic data, photographic images and physical modeling of volcanic plumes as a tool for monitoring the activity of Nevado del Ruiz Volcano. *Front. Earth Sci.* **6**, 162 (2018).
13. Ordoñez, M., Laverde, C. & Battaglia, M. The new lava dome growth of Nevado del Ruiz (2015–2021). *J. Volcanol. Geoth. Res.* **430**, 107626 (2022).
14. Sparks, J. Causes and consequences of pressurization in lava dome eruptions. *Earth Planet. Sci. Lett.* **150**, 177–189 (1997).
15. Stix, J., Calvache V, M. L. & Williams, S. N. Galeras volcano Colombia interdisciplinary study of a Decade Volcano. *J. Volcanol. Geotherm. Res.* **77**, 1–4 (1997).
16. Coppola, D. *et al.* Shallow magma convection evidenced by excess degassing and thermal radiation during the dome-forming Sabancaya eruption (2012–2020). *Bull. Volcanol.* **84**, 16 (2022).
17. Campion, R. *et al.* Breathing and coughing: The extraordinarily high degassing of popocatepetl volcano investigated with an SO₂ camera. *Front. Earth Sci.* **6**, 163 (2018).
18. Valade, S. *et al.* Lava dome cycles reveal rise and fall of magma column at Popocatepetl volcano. *Nat. Commun.* **14**, 3254 (2023).
19. Lages, J. *et al.* Volcanic gas emissions along the Colombian arc segment of the northern volcanic zone (CAS-NVZ): Implications for volcano monitoring and volatile budget of the Andean Volcanic Belt. *Geochem. Geophys. Geosyst.* **20**, 5057–5081 (2019).
20. Stix, J. Understanding fast and slow unrest at volcanoes and implications for eruption forecasting. *Front. Earth Sci.* **6**, 163 (2018).
21. Shinohara, H. A new technique to estimate volcanic gas composition: Plume measurements with a portable multi-sensor system. *J. Volcanol. Geoth. Res.* **143**, 319–333 (2005).
22. Aiuppa, A., Federico, C., Giudice, G. & Gurreri, S. Chemical mapping of a fumarolic field: La Fossa Crater, Vulcano Island (Aeolian Islands, Italy). *Geophys. Res. Lett.* **32**, 1–4 (2005).
23. Galle, B. *et al.* Network for observation of volcanic and atmospheric change (NOVAC)-a global network for volcanic gas monitoring: Network layout and instrument description. *J. Geophys. Res. Atmos.* **115**, 1–19 (2010).
24. Coppola, D., Laiolo, M., Cigolini, C., Delle Donne, D. & Ripepe, M. Enhanced volcanic hot-spot detection using MODIS IR data: Results from the MIROVA system. *Geol. Soc. Spec. Publ.* **426**, 181–205 (2016).
25. Aiuppa, A., Fischer, T. P., Plank, T., Robidoux, P. & Di Napoli, R. Along-arc, inter-arc and arc-to-arc variations in volcanic gas CO₂/S₂ ratios reveal dual source of carbon in arc volcanism. *Earth Sci. Rev.* **168**, 24–47 (2017).
26. Plank, T. The Chemical Composition of Subducting Sediments. In *Treatise on Geochemistry* 2nd edn, Vol. 4 607–629 (Elsevier, 2014).
27. Symonds, R. B., Gerlach, T. M. & Reed, M. H. Magmatic gas scrubbing: Implications for volcano monitoring. *J. Volcanol. Geoth. Res.* **108**, 303–341 (2001).
28. Casas, A. S. *et al.* SO₂ scrubbing during percolation through rhyolitic volcanic domes. *Geochim. Cosmochim. Acta* **257**, 150–162 (2019).
29. Aiuppa, A. *et al.* A model of degassing for Stromboli volcano. *Earth Planet. Sci. Lett.* **295**, 195–204 (2010).
30. Stix, J., Layne, G. D. & Williams, S. N. Mechanisms of degassing at Nevado del Ruiz volcano, Colombia. *J. Geol. Soc.* **160**, 507–521 (2003).
31. Gourgaud, A. & Thouret, J. C. Magma mixing and petrogenesis of the 13 November 1985 eruptive products at Nevado del Ruiz (Colombia). *J. Volcanol. Geoth. Res.* **41**, 79–96 (1990).

32. Melson, W. G. *et al.* Water contents, temperatures and diversity of the magmas of the catastrophic eruption of Nevado del Ruiz, Colombia, November 13, 1985. *J. Volcanol. Geoth. Res.* **41**, 97–126 (1990).
33. Sigurdsson, H., Carey, S., Palais, J. M. & Devine, J. Pre-eruption compositional gradients and mixing of andesite and dacite magma erupted from Nevado del Ruiz Volcano, Colombia in 1985. *J. Volcanol. Geoth. Res.* **41**, 127–151 (1990).
34. Plank, T. & Manning, C. E. Subducting carbon. *Nature* **574**, 343–352 (2019).
35. Wallace, P. J. Volatiles in subduction zone magmas: Concentrations and fluxes based on melt inclusion and volcanic gas data. *J. Volcanol. Geotherm. Res.* **140**, 217–240 (2005).
36. Moretti, R., Papale, P. & Ottonello, G. A model for the saturation of C-O-H-S fluids in silicate melts. *Geol. Soc. Spec. Publ.* **213**, 81–101 (2003).
37. Moretti, R. & Papale, P. On the oxidation state and volatile behavior in multicomponent gas-melt equilibria. *Chem. Geol.* **213**, 265–280 (2004).
38. Edmonds, M., Liu, E. J. & Cashman, K. V. Open-vent volcanoes fuelled by depth-integrated magma degassing. *Bull. Volcanol.* **84**, 28 (2022).
39. Coppola, D., Laiolo, M., Piscopo, D. & Cigolini, C. Rheological control on the radiant density of active lava flows and domes. *J. Volcanol. Geotherm. Res.* **249**, 39–48 (2013).
40. Shinohara, H. Excess degassing from volcanoes and its role on eruptive and intrusive activity. *Rev. Geophys.* **46**, (2008).
41. Lopez, T. *et al.* New insights into the magmatic-hydrothermal system and volatile budget of Lastarria volcano, Chile: Integrated results from the 2014 IAVCEI CCVG 12th volcanic gas workshop. *Geosphere* **14**, 983–1007 (2018).
42. Arellano, S. R. *et al.* Degassing patterns of Tungurahua volcano (Ecuador) during the 1999–2006 eruptive period, inferred from remote spectroscopic measurements of SO₂ emissions. *J. Volcanol. Geotherm. Res.* **176**, 151–162 (2008).
43. Williams, S. N. *et al.* Sulfur dioxide from Nevado del Ruiz volcano, Colombia: Total flux and isotopic constraints on its origin. *J. Volcanol. Geotherm. Res.* **42**, 53–68 (1990).
44. Krueger, A. J., Walter, L. S., Schnetzler, C. C. & Doiron, S. D. TOMS measurement of the sulfur dioxide emitted during the 1985 Nevado del Ruiz eruptions. *J. Volcanol. Geotherm. Res.* **41**, 7–15 (1990).
45. Stix, J. & de Moor, J. M. Understanding and forecasting phreatic eruptions driven by magmatic degassing. *Earth Planets Space* **70**, 83 (2018).
46. Giggenschbach, W. F. *et al.* The chemistry of fumarolic vapor and thermal-spring discharges from the Nevado del Ruiz volcanic-magmatic-hydrothermal system, Colombia. *J. Volcanol. Geotherm. Res.* **42**, 13–39 (1990).
47. Sturchio, N. C., Williams, S. N., Garcia, N. P. & Londono, A. C. The hydrothermal system of Nevado del Ruiz volcano Colombia. *Bull. Volcanol.* **50**, 399–412 (1988).
48. Lages, J. *et al.* Noble gas magmatic signature of the Andean Northern Volcanic Zone from fluid inclusions in minerals. *Chem. Geol.* **559**, 119966 (2021).
49. Naranjo, J. L., Sigurdsson, H., Carey, S. N. & Fritz, W. Eruption of the Nevado del Ruiz volcano, Colombia, on 13 November 1985: tephra fall and lahars. *Science* **233**, 961–963 (1986).
50. Aiuppa, A. *et al.* Volcanic CO₂ tracks the incubation period of basaltic paroxysms. *Sci. Adv.* **7**, eabh0191 (2021).
51. Lages, J. *et al.* First in-situ measurements of plume chemistry at Mount Garet volcano, island of Gaua (Vanuatu). *Appl. Sci.* **10**, 1–15 (2020).
52. Buck, A. L. New equations for computing vapor pressure and enhancement factor. *J. Appl. Meteorol. Climatol.* **20**, 1527–1532 (1981).
53. Tamburello, G. Ratiocalc: Software for processing data from multicomponent volcanic gas analyzers. *Comput. Geosci.* **82**, 63–67 (2015).
54. Galle, B. *et al.* A miniaturised ultraviolet spectrometer for remote sensing of SO₂ fluxes: A new tool for volcano surveillance. *J. Volcanol. Geotherm. Res.* **119**, 241–254 (2003).
55. Arellano, S. *et al.* Synoptic analysis of a decade of daily measurements of SO₂ emission in the troposphere from volcanoes of the global ground-based Network for observation of volcanic and atmospheric change. *Earth Syst. Sci. Data* **13**, 1167–1188 (2021).
56. Laiolo, M. *et al.* Shallow magma dynamics at open-vent volcanoes tracked by coupled thermal and SO₂ observations. *Earth Planet. Sci. Lett.* **594**, 117726 (2022).

Acknowledgements

We would like to thank Gloria Patricia Cortés and all the staff from the *Observatorio Vulcanológico y Sismológico de Manizales (Servicio Geológico Colombiano)* for their outstanding logistical and scientific support throughout this investigation. Initial funding for producing the NOVAC data set here presented was provided by the European Commission FP5 (DORSIVA project) and FP6 (NOVAC project), and the operation of the NOVAC network is funded through initiatives of the Volcano Observatories and support from the Volcano Disaster Assistance Program of the United States Geological Survey (USGS). S.A. acknowledges support from the Swedish National Space Agency (149/18). MIROVA is a collaborative project between the Universities of Turin and Florence (Italy) and is supported by the Italian Civil Protection Department. We acknowledge the LANCE-MODIS system (<http://lance-modis.eosdis.nasa.gov/>) for providing Level 1b MODIS data, as well as the crucial support from the Alfred P. Sloan Foundation (Deep Carbon Observatory/DECADE project; UniPa-CiW subcontract 10881-1262), the MIUR (under grant PRIN2017-2017LMNLAW) and the PE-PNRR project “Return” for funding a large portion of this investigation.

Author contributions

J.L., A.A. and Z.C. designed the study. J.L. drafted the manuscript with the help of all co-authors. J.L., Z.C., J.R., and G.T. processed the MultiGAS data. S.A., Z.C. and J.L. processed the SO₂ flux data. D.C., M.L. F.M. and C.L. contributed to the analysis of MODIS satellite data. M.B., J.P. and G.G. contributed to field work, instrument maintenance and data transmission. L.C. provided ash emission records from Nevado del Ruiz between 2018 and 2021. C.L. provided logistical and scientific support.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-024-51380-5>.

Correspondence and requests for materials should be addressed to J.L.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024