



Empirical Analysis of Global-Scale Natural Data and Analogue Seismotectonic Modelling Data to Unravel the Seismic Behaviour of the Subduction Megathrust

Francesca Funicello^{1*}, Fabio Corbi², Arnaud Heuret³, Claudia Piromallo⁴ and Matthias Rosenau⁵

¹Department of Science, University “Roma TRE”, Roma, Italy, ²Istituto di Geologia Ambientale Geoingegneria—CNR C/o Dipartimento Scienze della Terra, Sapienza Università di Roma, Roma, Italy, ³Université De Guyane / Géosciences Montpellier, Cayenne, France, ⁴Istituto Nazionale Di Geofisica e Vulcanologia, Roma, Italy, ⁵Helmholtz Centre Potsdam—GFZ German Research Centre for Geosciences, Potsdam, Germany

OPEN ACCESS

Edited by:

Debora Presti,
University of Messina, Italy

Reviewed by:

Magdala Tesauro,
Utrecht University, Netherlands
Luis E. Lara,
Servicio Nacional de Geología y
Minería de Chile (SERNAGEOMIN),
Chile

*Correspondence:

Francesca Funicello
francesca.funicello@uniroma3.it

Specialty section:

This article was submitted to
Structural Geology and Tectonics,
a section of the journal
Frontiers in Earth Science

Received: 28 August 2020

Accepted: 17 November 2020

Published: 17 December 2020

Citation:

Funicello F, Corbi F, Heuret A,
Piromallo C and Rosenau M (2020)
Empirical Analysis of Global-Scale
Natural Data and Analogue
Seismotectonic Modelling Data to
Unravel the Seismic Behaviour of the
Subduction Megathrust.
Front. Earth Sci. 8:600152.
doi: 10.3389/feart.2020.600152

Subduction megathrusts host the Earth’s greatest earthquakes as the 1960 Valdivia (M_w 9.5, Chile), the largest earthquake instrumentally recorded, and the recent 2004 Sumatra-Andaman (M_w 9.2, Indonesia), 2010 Maule (M_w 8.8, Chile), and 2011 Tohoku-Oki (M_w 9.1, Japan) earthquakes triggering devastating tsunamis and representing a major hazard to society. Unravelling the spatio-temporal pattern of these events is thus a key for seismic hazard assessment of subduction zones. This paper reviews the current state of knowledge of two research areas—empirical analysis of global-scale natural data and experimental data from an analogue seismotectonic modelling—devoted to study cause-effect relationships between subduction zone parameters and the megathrust seismogenic behavior. The combination of the two approaches overcomes the observational bias and inherent sampling limitations of geological processes (i.e., shortness of instrumental and historical data, decreasing completeness and resolution with time into the past) and allows drawing appropriately from multiple disciplines with the aim of highlighting the geodynamic conditions that may favor the occurrence of giant megathrust earthquakes.

Keywords: analogue seismotectonic modelling, analysis of global-scale natural data, mega-earthquakes, interplate seismicity, subduction megathrust

INTRODUCTION

Most of the global seismicity and the largest earthquakes occur at subduction zones where oceanic lithosphere is recycled as it sinks into the underlying mantle. The downgoing movement of the subducting plate and friction acting at the shallow part of the interface between the subducting and overriding plate (aka megathrust) cause shear stresses to accumulate slowly over centuries. When the frictional shear strength of the interface is overcome, these stresses are released episodically by a variety of seismic and aseismic slip modes, including “mega-earthquakes”: seismic megathrust rupture causing earthquakes with magnitudes (M_w) ≥ 8.5 . Stress accumulation and release occurs in cycles known as seismic cycles. In the last 15 years, a cluster of mega-earthquakes occurred at an

increased rate with respect to the last century (e.g., Ammon et al., 2010) often revealing unexpected characteristics (e.g., Stein and Okal 2011; Yue and Lay 2013) and resulting in catastrophic direct but also indirect (e.g., tsunamis) effects as in Sumatra (2004) (e.g., Subaraya et al., 2006), and Tohoku-Oki 2011 (e.g., Romano et al., 2012). The attention for these events has thus recently increased, raising interest on factors controlling their spatio-temporal pattern, including their potential maximum magnitude and return periods which are fundamental parameters in earthquake and tsunami hazard assessment. This is a challenging goal not only because each subduction zone has unique characteristics (e.g., Kopp, 2013) but also because the seismogenic portion of the megathrust is located offshore, below the sea-level (Heuret et al., 2011), and direct observation is restrained to shallow drilling (e.g., Japan Trench Fast Drilling Project; Chester et al., 2012) and the study of exhumed prototypes (e.g., Remitti et al., 2011). Moreover, the instrumental seismic record is limited to the last century, a very short time interval compared to the recurrence time of great earthquakes, which ranges from centuries to millennia (e.g., McCaffrey, 2008). In some cases, seismic records of megathrust ruptures can be extended back in time with paleoseismological investigations (e.g., Philibosian and Meltzner 2020), however such data typically lack the spatio-temporal completeness and resolution needed for rigorous statistical analysis.

One of the empirical research strategies that allow overcoming limited records consists in trading time with space at the global scale: given that different subduction zones are at individual stages of their seismic cycle, we may be able to infer the general seismogenic behavior of subduction zones by integrating information from worldwide seismicity (e.g., Schafer and Wenzel, 2019). The second approach, somehow complementary to the first one, is comparing the seismic character of individual megathrusts with the widest range of geological, geophysical and geodynamical parameters, with the aim of statistically identifying the combination of factors controlling subduction seismicity (e.g., Ruff and Kanamori, 1980; Peterson and Seno, 1984; Heuret et al., 2011; Schellart and Rawlinson, 2013; Brizzi et al., 2018).

Results from such a global-scale natural data analysis approach can be then tested against experimental and modelling data in order to validate inferred causal relationships by means of ad hoc parametric studies. Analytical (e.g., Ruff, 1992), numerical (e.g., van Dinther et al., 2013), physical (e.g., Den Hartog and Spiers 2013) and analog models (e.g., Rosenau et al., 2017) can reproduce a long series of seismic cycles and, by tuning ad hoc boundary conditions, they allow testing the role played on seismicity by the different parameters deduced from the analysis of natural data (e.g., Corbi et al., 2017b). Despite models are only simplified representations of the complexity of the natural prototype due to the unavoidable computational and experimental limitations, their results are inspiring for the interpretation of the limited and sparse natural data, thus enhancing our current understanding of subduction seismicity.

We focus our brief overview on the state of the art of the analysis from global-scale natural data and experimental data

from analogue seismotectonic modelling, highlighting the advantages of their use and the great potential for future developments in the study of the earthquake process that can be drawn from the close interaction and reciprocal feedback between the two methods. Very recently, also numerical methods have been developed to setup simulations which capture both the long-term and the short-term evolution of the subduction process. While numerical simulations proved particularly useful for testing hypothesis and mapping a wide parameter space, it is the emergent behavior of analogue models which makes them appealing for an empirical analysis as addressed in this review.

OBSERVATIONS AND INFERENCES FROM NATURAL DATA

There is a noteworthy variability in the maximum moment magnitude of earthquakes originated at the subduction megathrust during the instrumental era (e.g., Dixon and Moore, 2007 **Figure 1A**): some subduction zones—e.g., Alaska, Chile, Sumatra, NE Japan—experienced mega-earthquakes, while others—e.g., Tonga Mariana—did not (**Figure 1A**). What is unclear and still debated is if this dichotomy results only from the short observational time span and therefore all the worldwide subduction zones could host mega-earthquakes given a long enough observation period (e.g., McCaffrey, 2008) or if specific local conditions are responsible for the diversity of the interplate seismicity (e.g., Ruff and Kanamori, 1980). It is thus tempting to identify which geological/geophysical features, individually or in combination, possibly have a cause-effect relationship with megathrust seismicity (e.g., Pacheco et al., 1993; Heuret et al., 2011; Schellart and Rawlinson, 2013) and to quantify the likelihood for great earthquakes along worldwide subduction megathrusts (e.g., Marzocchi et al., 2016).

The “comparative subductology” was the intuition to disentangle the overall behavior of subduction by comparing a wide range of observables (i.e., subducting plate age and dip, convergent velocity, backarc tectonic style, characteristics of arc volcanism and seismicity, accretionary/erosional character of the margin) with seismicity characteristics of convergent margins (Uyeda, 1982) with the aim of deducing possible causal relationships (**Figure 1B**). This space-time trade-off approach allowed to overcome limitations due to uniqueness of local conditions (e.g., Suárez and Albin, 2009; Becker and Meier, 2010; Hayes and Furlong, 2010) and to the undersampling of the modern seismicity record.

Early on, two end-members subduction types — Mariana and Chilean — have been proposed, showing also significant difference in the degree of mechanical coupling (i.e., the ability to accumulate stresses) along the plate interface and in the consequent capability to host mega-earthquakes (Uyeda and Kanamori, 1979). The Chilean-type boundaries seemed more coupled and prone to trigger mega-earthquakes than the rather aseismic Marianas, with the different behaviors likely tuned by both different densities of the subducting plate and motions of the upper plate (Uyeda and Kanamori, 1979).




FIGURE 1 | (A) Map of the subduction interface seismicity (redrawn from Heuret et al., 2011). $M_w \geq 8.0$ subduction interface events are plotted by stars of different colours, representing the interval of occurrence of the events. The largest events are not homogeneously distributed in space and time along the worldwide subduction zones; **(B)** Correlation matrix used to show the dependence between multiple variables at a glance. The table contains the modulus of the Pearson's product moment correlation coefficient R of linear bivariate regressions between seismogenic zone parameters (t_{tot} : total seismic of subduction zones; t_{inter} : seismic rate due to interplate earthquakes; $Mmrr_{tot}$: total moment release rate of subduction zones; $Mmrr_{inter}$: moment release rate due to interplate events; $Mmax_{inter}$: maximum magnitude of interplate events; Cn : seismic coupling coefficient; $zmax_{inter}$: downdip limit of the subduction thrust fault; $zmin_{inter}$: updip limit of the subduction thrust fault) and other independent subduction parameters ($Tsed$: thickness of sediments at the trench; $UpPlate$: nature of the upper plate; thermal: thermal parameter; age: age of the subducting lithosphere at the trench; Vsn : normal component of the subduction velocity; $Vsupn$: normal component of the overriding plate velocity; Vtn : normal component of the trench velocity; $Vsubn$: normal component of the subducting plate velocity; Oc : convergence obliquity; Os : subduction obliquity; R : radius of bending of the slab) as attempt to unravel the overall behavior of subduction by comparing a wide range of observables (see Heuret et al., 2011 for additional details).

Based on this idea, Ruff and Kanamori (1980) looked at the maximum M_w recorded in the instrumental era and they found a striking correlation between the convergence rate and plate age. It was thus proposed that mega-earthquakes preferentially occur where a young lithosphere is converging fast with the overriding plate, allowing for a high mechanical plate coupling along the megathrust. This relationship has long been supported by studies involving larger, progressively updated and refined datasets (e.g., Peterson and Seno, 1984; Jarrard, 1986; Pacheco et al., 1993; Scholz and Campos, 1995). However, variations of coupling found within individual subduction zones (Scholz and Campos, 2012), the identification of different moment release rates in subduction zones sharing similar lithospheric ages and

speeds (Peterson and Seno, 1984) and, last but not least, updated data (e.g., Heuret et al., 2011; Stein and Okal, 2007; Stein and Okal, 2011) initiated skepticism in the acknowledgement of the Ruff and Kanamori (1980) model (e.g., McCaffrey, 1994) that was finally challenged by recent mega-earthquakes occurring in locations assumed to be weakly coupled (e.g., Wang, 2013). Among these, the Tohoku-Oki earthquake stands out not only for exceeding the regional maximum magnitude earthquake for that region according to Ruff and Kanamori (1980) model, but also as case of extreme violation of commonly accepted scaling laws (e.g. Okal, 2015).

A new phase of the comparative subductology involved not only the use of large dataset (e.g., Heuret et al., 2012; Schellart and

Rawlinson 2013) but also of more sophisticated statistical analysis to treat them (e.g., Heuret et al., 2012; Marzocchi et al., 2016; Brizzi et al., 2018; Schafer and Wenzel, 2019). Cause-effect relationships between geodynamical and seismological parameters, have inspired physical models of seismically active convergent margins highlighting the possible role of numerous factors in triggering mega-earthquakes: trench sediments (Kostoglodov, 1988; Ruff, 1989; Heuret et al., 2011; Heuret et al., 2012; Scholl et al., 2015; Seno, 2017; Brizzi et al., 2018), plate roughness (Morgan et al., 2008; Kopp, 2013; Wang and Bilek, 2014; Bassett and Watts, 2015; Lallemand et al., 2018; van Rijnsingen et al., 2018) upper plate strain (Conrad et al., 2004; Heuret et al., 2011; Heuret et al., 2012; Schellart and Rawlinson, 2013), plate age (Nishikawa and Ide, 2014), forearc structure (Song and Simons, 2003; Wells et al., 2003; Fuller et al., 2006; Rosenau and Oncken, 2009), along-strike (Schellart and Rawlinson, 2013) and along-dip (Bletery et al., 2016) curvature of subduction zones, slab dip (Schafer and Wenzel, 2019), downdip width of the seismogenic zone (e.g., Kelleher et al., 1974; Pacheco et al., 1993; Hayes et al., 2012; Schellart and Rawlinson, 2013; Schafer and Wenzel, 2019) and propensity to produce large rupture in the trench-parallel direction (Bilek and Lay, 1999; Brizzi et al., 2018). The proposed models highlight ingredients able to produce a homogeneous and strong subduction megathrust allowing ruptures to easily propagate along-strike and, thus, to grow into mega-earthquakes.

Such a physical understanding may re-evaluate traditional probabilistic seismic-hazard assessment, suggesting that, irrespective of the geodynamic context, many subduction zones (Berryman et al., 2015) if not all (McCaffrey, 2008; Rong et al., 2014) may have the potential of producing mega-earthquakes (McCaffrey, 2008; Rong et al., 2014) but they are likely characterized by a difference in the average rate of occurrence of these events (Marzocchi et al., 2016).


INSIGHTS FROM ANALOG MODELLING OF SUBDUCTION MEGATHRUST SEISMIC CYCLES

To overcome limitations of natural records and test cause-effect relationships suggested by global-scale data analysis we promote an analog modelling approach. Analog models are down-scaled and simplified versions of natural subduction zones used in experiments that allow reproducing the essence of a variety of geological phenomena, in convenient spatial and temporal scales (e.g., Hubbert, 1937; Funicello and Corbi, 2017). Despite analog models might appear oversimplified with respect to high performance computational modelling strategies anticipated in the introduction (for example, no efficient control on temperature and water content has been implemented so far), they have the advantage of evolving spontaneously on the basis of the physics of the process, which can be easy to tune and to interpret. Moreover, analog models are intrinsically three-dimensional and occur in a space-time continuum, both keys in allowing across-scale processes to emerge, as e.g. the growth of mega-earthquakes (e.g., Brizzi et al., 2018). The recent advances

in characterization of rheological (e.g., Di Giuseppe et al., 2014; Rudolf et al., 2016) and frictional properties (Klinkmüller et al., 2016) of analog materials and the improvement in monitoring techniques (Rudolf et al., 2019) supported the development of so-called “seismotectonic” analog models (Rosenau et al., 2017). Those models reproduce the elastic loading and sudden release in a cyclic fashion similarly to the seismic cycle of subduction megathrusts (**Figure 2**). All seismotectonic analog models developed so far share the common characteristics listed below:

- *Kinematic boundary condition*: deformation of the overriding plate is imposed by underthrusting of the subducting plate at a constant rate.
- *Wedge geometry*: elastic stresses are stored into the analog overriding plate which might be elastic or elastoplastic while the subducting plate is assumed being rigid; the initial wedge surface and the analog megathrust are flat (i.e., with no downdip- and along trench curvature).
- *Rate-state friction*: the analog megathrust includes both (aseismic) velocity-strengthening and (seismogenic) velocity-weakening frictional regions.
- *Spontaneous earthquake nucleation*: slip episodes occur spontaneously when elastic stress overcomes the frictional strength of the megathrust locally (i.e., there is no external triggering).
- *Laboratory geodetic monitoring*: models are monitored via cross-correlation of images of the top or side surfaces or the interior of the model (e.g., with the Particle Image Velocimetry, PIV method; Svein 2004). Applied to the top surface this technique is equivalent to having a dense network of continuous GPS stations (hence the name “laboratory geodesy models”) which allow inverting for relevant earthquake source parameters (e.g., Rosenau et al., 2019; Corbi et al., 2013; Kosari et al., 2020).

The first articles about analog modelling of megathrust seismic cycling focused on reporting rheological and frictional properties of analog materials (Corbi et al., 2011) and their implementation into a scale model (Rosenau et al., 2009). The experimental behavior was composed of the archetypical pattern of alternating relatively longer periods of low, landward deformation rates with sudden, higher, trenchward velocities reversals as in the seismic cycle of subduction megathrusts (Rosenau et al., 2009; Corbi et al., 2013; **Figure 2** panels c–f). Follow-up studies focused on applications demonstrating the capabilities of analog models in studying a variety of earthquake phenomena, including recurrence pattern (Rosenau and Oncken, 2009; Rosenau et al., 2019) and the moment-duration proportionality (Corbi et al., 2013). Analog models have been used for investigating extrinsic parameter control, for example how the width of the seismogenic zone (Corbi et al., 2017a), the upper plate rheology (Brizzi et al., 2016), the subduction velocity (Corbi et al., 2017a) or the subduction megathrust roughness (van Rijnsingen et al., 2019) influence the seismic behavior. Analog models helped in constraining the intrinsic variability of earthquake ruptures and associated tsunamis, validating empirical tsunami forecast



models (Rosenau et al., 2010). Analog models have been also used for identifying the geometrical and mechanical conditions leading to synchronization of earthquakes (Corbi et al., 2017b; Rosenau et al., 2019). The process of earthquake synchronization is one of the key aspects in seismogenesis and is considered responsible for the origin of mega-earthquakes unzipping large parts of a subduction zone. Nowadays, analog models are also used for testing earthquake predictability (e.g., Corbi et al., 2019) and for identifying optimal geodetic configurations for monitoring convergent margins (Corbi et al., 2020; Kosari et al., 2020).

CHALLENGES AND FUTURE DIRECTIONS

Over the last decades, a wide range of studies have shed light on the seismogenic behavior of subduction megathrusts. This short review has recapped the contributions provided by the comparative analysis of global-scale natural data and analogue modelling.

Forecasting which subduction zone will host the next mega-earthquake and its timing remains a major challenge. The efforts to highlight which are the conditions enhancing the “productivity” of mega-earthquakes have recognized the potential role of several subduction-related parameters, but no consensus has been reached so far. For instance, empirical results point out at different parameters such as plates curvature (Bletery et al., 2016), roughness (e.g., van Rijnsingen et al., 2019) or trench-parallel length of the subduction zone (Brizzi et al., 2018) and amount of trench sediments (Brizzi et al., 2020). Convergent margins are complex tectonic settings, whose diversity results from the interplay of several parameters operating at different spatial and temporal scales and whose relative importance is thus hard to assess (e.g., Figure 1B). Moreover, the understanding of subduction megathrusts is biased by the short instrumental seismic record and by the lack of completeness and of a homogeneous resolution of the observables characterizing subduction zones.

Is it thus really feasible to unravel the seismic behavior of the subduction megathrust or is it an impossible mission?

The continuous scientific and technological advancements allow detecting transient deformation signals occurring on short times-scales (fractions of the seismic cycle) along convergent margins and sampling the megathrust seismic cycle with progressively denser and higher quality data. For example, even if the underlying source process still has to be identified, both the 2010 Maule (Chile) and 2011 Tohoku-Oki (Japan) earthquakes have been preceded by potentially diagnostic geodetic transients lasting several months and spanning thousands of kilometers (Bedford et al., 2020). Satellite measurements (e.g., InSAR studies; Jolivet et al., 2020), dense geodetic (e.g., Michel et al., 2018) and seismic (e.g., Brodsky and Lay, 2014) monitoring - both onshore and offshore - is able to record a wide range of deformation signal frequencies (i.e., from earthquakes s.s. to low frequency earthquakes and tremors) and the improvement of the quantity, quality and accessibility of databases of convergent margin observables (e.g., geodesy.unr.edu/NGLStationPages/gpsnetmap/GPSNetMap.html; geofon.gfz-potsdam.de; www-solid.eps.s.u-tokyo.ac.jp/~sloweq) yield a promising perspective to successfully elucidate the spatio-temporal pattern of earthquake occurrence along the worldwide subduction zones and to mitigate hazard related to mega-earthquakes and tsunamis.

Another support to better understand subduction earthquakes and to advance in the forecast of earthquakes is coming from recent improvements in data analysis methods. Machine learning is an application of artificial intelligence able to automatically discover hidden patterns learning directly from analysis of large datasets, without being explicitly programmed (e.g., Bergen et al., 2019; Ren et al., 2020). This approach already succeeded for the prediction of timing (Rouet-Leduc et al., 2018), timing and locations (Corbi et al., 2019) and timing and duration (Hulbert et al., 2019) of slip episodes in laboratory experiments mimicking the basic physics of the seismic cycle.

The future is challenging also on the analogue modelling side, where the use of higher resolution and faster cameras and of optimized procedures for calculating surface velocities (Rudolf et al., 2019) permits the model monitoring with increasing detail. Thus, the detection threshold of analog earthquakes as well as the temporal resolution of the nucleation and rupture process are progressively improved. Laboratory geodetic methods will be accompanied by laboratory seismology using micro-electromechanical sensors (MEMS) measuring model accelerations with kHz sampling rates, corresponding to the frequency range of interest for earthquakes. This will soon put us in a position to detect and localize analog earthquakes which are too small to produce a surface deformation signal measurable with optical methods. Apart from such kinematic monitoring methods, dynamic monitoring of (boundary) stresses (Ritter et al., 2018) will help in detecting events but also in

understanding the relationship between stress and strain during seismic cycles. Moreover, the joint use of analogue and numerical modelling is providing successful and promising results to shed light on the behavior of the subduction megathrust (Corbi et al., 2013, Corbi et al., 2017a; van Dinther et al., 2013). The combination of these two complementary approaches allows to valorize the respective strengths: analog models are physically self-consistent, and stresses/strain evolve spontaneously in response to the applied boundary conditions, while numerical models are more adaptable and effective for parametric studies.

With respect to great subduction megathrust earthquakes, our current understanding is that they show a complex but in parts deterministic pattern. Whether (mega-) earthquakes are predictable or not can be thus answered with some hopefulness in the future. Since the seismic behavior of the subduction megathrust depends on a wide combination of parameters, interdisciplinary research needs to be strengthened integrating geological, geophysical, and geodynamical constraints and contextualizing them in a physics-based modelling framework allowing to analyze and forecast mega-earthquake triggering processes in subduction zones.

Multidisciplinary studies are always challenging, but they represent the key for a substantial advancement in the understanding of this scientifically intriguing and socially hazardous issue. In this frame, collecting new data, sharing them on open platforms, networking resources and events are fundamental steps for exchanging ideas and fostering discussion towards the comprehension of the processes controlling the megathrust seismic behavior.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

ACKNOWLEDGMENTS

The authors acknowledge the Editor, DP, and two reviewers for their suggestions which help to improve the final version of the manuscript. The Grant to Department of Science, Roma Tre University (MIUR-Italy Dipartimenti di Eccellenza, ARTICOLO 1, COMMI 314–337, LEGGE 232/2016) is gratefully acknowledged by FF. FC thanks his funding from PRIME program of the German Academic Exchange Service (DAAD) with funds from the German Federal Ministry of Education and Research (BMBF). MR has been supported by the Deutsche Forschungsgemeinschaft (DFG) through grant CRC 1114 “Scaling Cascades in Complex Systems,” Project Number 235221301, Project B01.

REFERENCES

- Ammon, C. J., Lay, T., and Simpson, D. W. (2010). Great earthquakes and global seismic networks. *Seismol. Res. Lett.* 6 (81), 965–971. doi:10.1785/gssrl.81.6.965
- Bassett, D., and Watts, A. (2015). Gravity anomalies, crustal structure, and seismicity at subduction zones: 1. seafloor roughness and subducting relief. *Geochem. Geophys. Geosyst.* 16, 1508–1540. doi:10.1002/2014GC005684
- Becker, D., and Meier, T. (2010). Seismic dip deficit in the southwestern forearc of the hellenic subduction zone. *Bull. Seismol. Soc. Am.* 100, 325–342. doi:10.1785/0120090156
- Bedford, J. R., Moreno, M., Deng, Z., Oncken, O., Schurr, B., John, T., et al. (2020). Months-long thousand-kilometre-scale wobbling before great subduction earthquakes. *Nature*. 580 (7805), 628–635. doi:10.1038/s41586-020-2212-1
- Bergen, K. J., Johnson, P. A., De Hoop, M. V., and Beroza, G. C. (2019). Machine learning for data-driven discovery in solid Earth geoscience. *Science*. 363 (6433). doi:10.1126/science.aau0323

- Berryman, K., Wallace, L., Hayes, G., Bird, P., Wang, K., Basili, R., et al. (2015). The GEM faulted earth subduction interface characterization project. Version 2.0. <http://www.nexus.globalquakemodel.org/gem-faulted-earth/posts>.
- Bilek, S. L., and Lay, T. (1999). Rigidity variations with depth along interplate megathrust faults in subduction zones. *Nature*. 446 (July), 443–446. doi:10.1038/22739
- Bletery, Q., Thomas, A. M., Rempel, A. W., Karlstrom, L., Sladen, A., and De Barros, L. (2016). Mega-earthquakes rupture flat megathrusts. *Science*. 354, 1027–1031. doi:10.1126/science.aag0482
- Brizzi, S., Funicello, F., Corbi, F., Di Giuseppe, E., and Mojoli, G. (2016). Salt matters: how salt affects the rheological and physical properties of gelatine for analogue modelling. *Tectonophysics*. 679, 88–101. doi:10.1016/j.tecto.2016.04.021
- Brizzi, S., Sandri, L., Funicello, F., Corbi, F., Piromallo, C., and Heuret, A. (2018). Multivariate statistical analysis to investigate the subduction zone parameters favoring the occurrence of giant megathrust earthquakes. *Tectonophysics*. 728–729, 92–103. doi:10.1016/j.tecto.2018.01.027
- Brizzi, S., Zelst, I., Funicello, F., Corbi, F., and van Dinther, Y. (2020). How sediment thickness influences subduction dynamics and seismicity. *J. Geophys. Res. Solid Earth*. 125 (8), 1–19. doi:10.1029/2019jb018964
- Brodsky, E. E., and Lay, T. (2014). Recognizing foreshocks from the 1 april 2014 Chile earthquake. *Science*. 344 (May), 700–702. doi:10.1126/science.1255202
- Chester, F. M., Mori, J. J., Toczko, S., Eguchi, N., Kido, Y., et al. (2012). *Integrated ocean drilling program expedition 343/343T preliminary report, Japan Trench Fast Drilling Project (JFAST) Prelim. Rep. 343: Integrated Ocean Drilling Project*.
- Conrad, C. P., Bilek, S., and Lihtgow-Bertelloni, C. (2004). Great earthquakes and slab pull: interaction between seismic coupling and plate-slab coupling. *Earth Planet Sci. Lett.* 218, 109–122. doi:10.1016/s0012-821x(03)00643-5
- Corbi, F., Funicello, F., Faccenna, C., Ranalli, G., and Heuret, A. (2011). Seismic variability of subduction thrust faults: insights from laboratory models. *J. Geophys. Res.* 116 (B6), 1–14. doi:10.1029/2010JB007993
- Corbi, F., Funicello, F., Moroni, M., van Dinther, Y., Mai, P. M., Dalguer, L. A., et al. (2013). The seismic cycle at subduction thrusts: 1. Insights from laboratory models. *J. Geophys. Res. Solid Earth*. 118, 1–19. doi:10.1002/2016GL072415
- Corbi, F., Funicello, F., Brizzi, S., Lallemand, S., and Rosenau, M. (2017a). Control of asperities size and spacing on seismic behavior of subduction megathrusts. *Geophys. Res. Lett.* 44, 8227–8235. doi:10.1002/2017GL074182
- Corbi, F., Herrendörfer, R., Funicello, F., and van Dinther, Y. (2017b). Controls of seismic zone width and subduction velocity on interplate seismicity: insights from analog and numerical models. *Geophys. Res. Lett.* 44, 6082–6091. doi:10.1002/2016GL072415
- Corbi, F., Sandri, L., Bedford, J., Funicello, F., Brizzi, S., Rosenau, M., et al. (2019). Machine learning can predict the timing and size of analog earthquakes. *Geophys. Res. Lett.* 46, 1303–1311. doi:10.1029/2018GL081251
- Corbi, F., Bedford, J., Sandri, L., Funicello, F., Gualandi, A., and Rosenau, M. (2020). Predicting imminence of analog megathrust earthquakes with machine learning: implications for monitoring subduction zones. *Geophys. Res. Lett.* 47 (7), e2019GL086615. doi:10.1029/2019GL086615
- Den Hartog, S. A. M., and Spiers, C. J. (2013). Influence of subduction zone conditions and gouge composition on frictional slip stability of megathrust faults. *Tectonophysics*. 600, 75–90. doi:10.1016/j.tecto.2012.11.006
- Di Giuseppe, E., CorbiFunicello, F. F., Massmeyer, A., Santimano, T. N., Rosenau, M., and Davaille, A. (2014). Tectonophysics Characterization of Carbopol[®] hydrogel rheology for experimental tectonics and geodynamics. *Tectonophysics*. [Epub ahead of print]. doi:10.1016/j.tecto.2014.12.005
- Dixon, T., and Moore, C. (2007). *The seismic zone of subduction thrust faults*. New York: Columbia Univ. Press, 123–146.
- Fuller, W., Willett, S. W., and Brandon, M. T. (2006). Formation of forearc basins and their influence on subduction zone earthquakes. *Geology*. 34 (2), 65–68. doi:10.1130/G21828.1
- Funicello, F., and Corbi, F. (2017). “Experimental tectonics: convergent margins from a lithosphere–mantle perspective,” in *Reference module in Earth Systems and environmental sciences*. (Elsevier). doi:10.1016/B978-0-12-409548-9.10019-3
- Hayes, G. P., and Furlong, K. P. (2010). Quantifying potential tsunami hazard in the puysegur subduction zone, south of new zealand. *Geophys. J. Int.* 183, 1512–1524. doi:10.1111/j.1365-246x.2010.04808.x
- Hayes, G. P., Wald, D. J., and Johnson, R. L. (2012). Slab1.0: a three-dimensional model of global subduction zone geometries. *J. Geophys. Res. Solid Earth*. 117, B01302. doi:10.1029/2011JB008524. b01302
- Heuret, A., Conrad, C. P., Funicello, F., Lallemand, S., and Sandri, L. (2012). Relation between subduction megathrust earthquakes, trench sediment thickness and upper plate strain. *Geophys. Res. Lett.* 39, 1–6. doi:10.1029/2011GL050712
- Heuret, A., Lallemand, S., Funicello, F., Piromallo, C., and Faccenna, C. (2011). Physical characteristics of subduction interface type seismogenic zones revisited. *Geochem. Geophys. Geosyst.* 12, 1–26. doi:10.1029/2010GC003230
- Hubbert, M. K. (1937). Theory of scale models as applied to the study of geological structures. *Geol. Soc. Am. Bull.* 48, 1459–1520. doi:10.1130/gsab-48-1459
- Hulbert, C., Rouet-Leduc, B., Johnson, P. A., Ren, C. X., Rivière, J., Bolton, D. C., et al. (2019). Similarity of fast and slow earth- quakes illuminated by machine learning. *Nat. Geosci.* 12 (1), 69–74. doi:10.1038/s41561-018-0272-8
- Jarrard, R. D. (1986). Relations among subduction parameters. *Rev. Geophys.* 24, 217–284. doi:10.1029/RG024i002p00217
- Jolivet, R., Simons, M., Duputel, Z., Olive, J.-A., Bhat, H. S., and Bletery, Q. (2020). Interseismic coupling drives long-term uplift in northern chile. *Geophys. Res. Lett.* 47, e2019GL085377. doi:10.1029/2019GL085377
- Kelleher, J., Savino, J., Rowlett, H., and McCann, W. (1974). Why and where great thrust earthquakes occur along island arcs. *J. Geophys. Res.* 79, 4889–4899. doi:10.1029/JB079i032p048
- Klinkmüller, M., Schreurs, G., Rosenau, M., and Kemnitz, H. (2016). Properties of granular analogue materials: a community wide survey. *Tectonophysics*. 666, 23–38. doi:10.1016/j.tecto.2016.01.017
- Kopp, H. (2013). Invited review paper: the control of subduction zone structural complexity and geometry on margin segmentation and seismicity. *Tectonophysics*. 589, 1–16. doi:10.1016/j.tecto.2012.12.037
- Kosari, E., Rosenau, M., Bedford, J., Rudolf, M., and Oncken, O. (2020). On the relationship between offshore geotectonic coverage and slip model uncertainty: analog megathrust earthquake case studies. *Geophys. Res. Lett.* 47 (15), e2020GL088266. doi:10.1029/2020gl088266
- Kostoglodov, V. (1988). Sediment subduction: a probable key for seismicity and tectonics at active plate boundaries. *Geophys. J.* 94, 65–72.
- Lallemand, S., Peyret, M., van Rijsingen, E., Arcay, D., and Heuret, A. (2018). Roughness characteristics of oceanic seafloor prior to sub- duction in relation to the seismogenic potential of subduction zones. *Geochem. Geophys. Geosyst.* 19, 1–26. doi:10.1029/2018GC007434
- Marzocchi, W., Sandri, L., Heuret, A., and Funicello, F. (2016). Where giant earthquakes may come. *J. Geophys. Res. Solid Earth*. 121, 7322–7336. doi:10.1002/2016JB013054
- McCaffrey, R. (1994). Dependence of earthquake size distributions on convergence rates at subduction zones. *Geophys. Res. Lett.* 21, 2327–2330. doi:10.1029/94gl02153
- McCaffrey, R. (2008). Global frequency of magnitude 9 earthquakes. *Geology*. 36, 263–266. doi:10.1130/G24402A.1
- Michel, S., Gualandi, A., and Avouac, J.-P. (2018). Interseismic coupling and slow slip events on the cascadia megathrust. *Pure Appl. Geophys.* 176 (10), 4641–4642. doi:10.1007/s00024-018-1991-x
- Morgan, E. C., McAdoo, B. G., and Baise, L. G. (2008). Quantifying geomorphology associated with large subduction zone earthquakes. *Basin Res.* 20, 531–542. doi:10.1111/j.1365-2117.2008.00368.x
- Nishikawa, T., and Ide, S. (2014). Earthquake size distribution in subduction zones linked to slab buoyancy. *Nat. Geosci.* 7 (12), 904–908. doi:10.1038/ngeo2279
- Okal, E. A. (2015). The quest for wisdom: lessons from 17 tsunamis, 2004–2014. *Phil. Trans. R. Soc. A* 373, 20140370. doi:10.1098/rsta.2014.0370
- Pacheco, J. F., Sykes, L. R., and Scholz, C. H. (1993). Nature of seismic coupling along simple plate boundaries of the subduction type. *J. Geophys. Res.* 98, 14133. doi:10.1029/93JB00349
- Peterson, E. T., and Seno, T. (1984). Factors affecting seismic moment release rates in subduction zones. *J. Geophys. Res.* 89, 10233–10248. doi:10.1029/jb089ib12p10233
- Philibosian, B., and Meltzner, A. J. (2020). Segmentation and supercycles: a catalog of earthquake rupture patterns from the Sumatran Sunda Megathrust and other well-studied faults worldwide. *Quat. Sci. Rev.* 241, 106390. doi:10.1016/j.quascirev.2020.106390
- Remitti, F., Vannucchi, P., Bettelli, G., Fantoni, L., Panini, F., and Vescovi, P. (2011). Tectonic and sedimentary evolution of the frontal part of an ancient subduction complex at the transition from accretion to erosion: the case of the

- Ligurian wedge of the northern apennines, italy. *Geol. Soc. Am. Bull.* 123 (1–2), 51–70. doi:10.1130/B30065.1
- Ren, C., Hulbert, C., Johnson, P., and Rouet-Leduc, B. (2020). Machine learning and fault rupture: review. Preprint repository name [Preprint]. Available at: <https://osf.io/g2dt8/>. doi:10.31223/osf.io/xdmkq
- Ritter, M. C., Santimano, T., Rosenau, M., Leever, K., and Oncken, O. (2018). Sandbox rheometry: co-evolution of stress and strain in riedel- and critical wedge-experiments. *Tectonophysics*. 722 (2), 400–409. doi:10.1016/j.tecto.2017.11.018
- Romano, F., Piatanesi, A., Lorito, S., D'Agostino, N., Hirata, K., Atzori, S., et al. (2012). Clues from joint inversion of tsunami and geodetic data of the 2011 Tohoku-oki earthquake. *Sci. Rep.* 2, 385. doi:10.1038/srep00385
- Rong, Y., Jackson, D. D., Magistrale, H., and Goldfinger, C. (2014). Magnitude limits of subduction zone earthquakes. *Bull. Seismol. Soc. Am.* 104, 2359–2377. doi:10.1785/0120130287
- Rosenau, M., Lohrmann, J., and Oncken, O. (2009). Shocks in a box: an analogue model of subduction earthquake cycles with application to seismotectonic forearc evolution. *J. Geophys. Res.* 114 (B1), 1–20. doi:10.1029/2008JB005665
- Rosenau, M., Oppelt, A., Kemnitz, H., and Oncken, O. (2008). Analogue earthquakes: friction experiments with rice and implications for the seismic cycle and earthquake nucleation. *Boll. Geofis.* 49, 104–108.
- Rosenau, M., Nerlich, R., Brune, S., and Oncken, O. (2010). Experimental insights into the scaling and variability of local tsunamis triggered by giant subduction megathrust earthquakes. *J. Geophys. Res.* 115, B09314. doi:10.1029/2009jb007100
- Rosenau, M., Corbi, F., and Dominguez, S. (2017). Analogue earthquakes and seismic cycles: experimental modelling across timescales. *Solid Earth*. 8 (3), 597–635. doi:10.5194/se-8-597-2017
- Rosenau, M., Horenko, I., Corbi, F., Rudolf, M., Kornhuber, R., and Oncken, O. (2019). Synchronization of great subduction megathrust earthquakes: insights from scale model analysis. *J. Geophys. Res. Solid Earth*. 124 (4), 3646–3661. doi:10.1029/2018JB016597
- Rosenau, M., and Oncken, O. (2009). Fore-arc deformation controls frequency-size distribution of megathrust earthquakes in subduction zones. *J. Geophys. Res. Solid Earth*. 114, 12. doi:10.1029/2009JB006359
- Rouet-Leduc, B., Hulbert, C., Bolton, D. C., Ren, C. X., Riviere, J., Marone, C., et al. (2018). Estimating fault friction from seismic signals in the laboratory. *Geophys. Res. Lett.* 45, 1321–1329. doi:10.1002/2017GL076708
- Rudolf, M. (2019). Velocity dependent rheologies in seismotectonic scale modelling: characterization and Implementation in a new analogue modelling scheme. PhD thesis. Berlin: Freie Universität, 135.
- Rudolf, M., Boutelier, D., Rosenau, M., and Schreurs, G., and Oncken, O. (2016). Rheological benchmark of silicone oils used for analog modeling of short- and long-term lithospheric deformation. *Tectonophysics*. 684, 12–22. doi:10.1016/j.tecto.2015.11.028
- Rudolf, M., Rosenau, M., Ziegenhagen, T., Ludwikowski, V., Schucht, T., Nagel, H., et al. (2019). Smart speed imaging in digital image correlation: application to seismotectonic scale modeling. *Front. Earth Sci.* 6, 248. doi:10.3389/feart.2018.00248
- Ruff, L. J. (1989). Do trench sediments affect great earthquake occurrence in subduction zones?. *Pure Appl. Geophys.* 129, 263–282. doi:10.1007/bf00874629
- Ruff, L. J. (1992). Asperity distributions and large earthquake occurrence in subduction zones. *Tectonophysics*. 211 (1–4), 61–83. doi:10.1016/0040-1951(92)90051-7
- Ruff, L., and Kanamori, H. (1980). Seismicity and the subduction process. *Phys. Earth Planet. In.* 23, 240–252. doi:10.1016/0031-9201(80)90117-X
- Schäfer, A. M., and Wenzel, F. (2019). Global megathrust earthquake hazard—maximum magnitude assessment using multi-variate machine learning. *Front. Earth Sci.* 7 (June), 1–19. doi:10.3389/feart.2019.00136
- Schellart, W. P., and Rawlinson, N. (2013). Global correlations between maximum magnitudes of subduction zone interface thrust earthquakes and physical parameters of subduction zones. *Phys. Earth Planet. In.* 225, 41–67. doi:10.1016/j.pepi. 2013.10.001
- Scholl, D. W., Kirby, S. H., von Huene, R., Ryan, H., Wells, R. E., and Geist, E. L. (2015). Great (\geq Mw8.0) megathrust earthquakes and the subduction of excess sediment and bathymetrically smooth seafloor. *Geosphere*. 11, 236–265. doi:10.1130/GES01079.1
- Scholz, C. H., and Campos, J. (1995). On the mechanism of seismic decoupling and back arc spreading at subduction zones. *J. Geophys. Res.* 100, 22103. doi:10.1029/95JB01869
- Scholz, C. H., and Campos, J. (2012). The seismic coupling of subduction zones revisited. *J. Geophys. Res.* 117, B05310. doi:10.1029/2011jb009003
- Seno, T. (2017). Subducted sediment thickness and Mw 9 earthquakes. *J. Geophys. Res. Solid Earth*. 122, 470–491. doi:10.1002/2016JB013048
- Song, T. R. A., and Simons, M. (2003). Large trench-parallel gravity variations predict seismogenic behaviour in subduction zones. *Science*. 301, 630–633. doi:10.1126/science.1085557
- Stein, S., and Okal, E. (2011). The size of the 2011 Tohoku earthquake need not have been a surprise. *EOS Trans. Am. Geophys. Union*. 92, 227–228. doi:10.1029/2011EO270005
- Stein, S., and Okal, E. (2007). Ultralong period seismic study of the december 2004 indian ocean earthquake and implications for regional tectonics and the subduction process. *Seismol. Soc. Am. Bull.* 97, S279–S295. doi:10.1785/0120050617
- Suarez, G., and Albin, P. (2009). Evidence for great tsunamigenic earthquakes along the mexican subduction zone. *Bull. Seismol. Soc. Am.* 99, 892–896. doi:10.1785/0120080201
- Subarya, C., Chlieh, M., Prawirodirdjo, L., Avouac, J. P., Bock, Y., Sieh, K., et al. (2006). Plate-boundary deformation associated with the great Sumatra-Andaman earthquake. *Nature*. 440, 46–51. doi:10.1038/nature04522
- Sveen, J. K. (2004). An introduction to MatPIV v.1.6.1: Department of Mathematics University of Oslo.
- Uyeda, S., and Kanamori, H. (1979). Back-arc opening and the mode of subduction. *J. Geophys. Res.* 84, 1049. doi:10.1029/JB084iB03p01049
- Uyeda, S. (1982). Subduction zones: an introduction to comparative subductology. *Tectonophysics*. 81 (3–4), 133–159. doi:10.1016/0040-1951(82)90126-3
- van Dinther, Y., Gerya, T. V., Dalguer, L. A., Corbi, F., Funicello, F., and Mai, P. M. (2013). The seismic cycle at subduction thrusts: 2. Dynamic implications of geodynamic simulations validated with laboratory models. *J. Geophys. Res. Solid Earth*. 118, 1502–1525. doi:10.1029/2012JB009479
- van Rijnsingen, E., Lallemand, S., Peyret, M., Arcay, D., Heuret, A., Funicello, F., et al. (2018). How subduction interface roughness influences the occurrence of large interplate earthquakes. *Geochem. Geophys. Geosyst.* 19, 1–29. doi:10.1029/2018GC007618
- van Rijnsingen, E., Funicello, F., Corbi, F., and Lallemand, S. (2019). Rough subducting seafloor reduces interseismic coupling and mega-earthquake occurrence: insights from analogue models. *Geophys. Res. Lett.* 46 (6), 3124–3132. doi:10.1029/2018GL081272
- Wang, K., and Bilek, S. L. (2014). Invited review paper: fault creep caused by subduction of rough seafloor relief. *Tectonophysics*. 610, 1–24. doi:10.1016/j.tecto.2013.11.024
- Wang, K. (2013). Megathrust surprises. *Nat. Geosci.* 6, 11–12. doi:10.1038/ngeo1682
- Wells, R. E., Blakely, R. J., Sugiyama, Y., Scholl, D. W., and Dinterman, P. A. (2003). Basin-centered asperities in great subduction zone earthquakes: a link between slip, subsidence, and subduction erosion?. *J. Geophys. Res.* 108, 2507. doi:10.1029/2002JB002072
- Yue, H., and Lay, T. (2013). Source rupture models for the Mw 9.0 2011 Tohoku earthquake from joint inversions of high-rate geodetic and seismic data. *Bull. Seismol. Soc. Am.* 103, 1242–1255. doi:10.1785/0120120119

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Funicello, Corbi, Heuret, Piromallo and Rosenau. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.