

On computing viscoelastic Love numbers for general planetary models: the ALMA³ code

D. Melini¹, C. Saliby² and G. Spada³

¹ *Istituto Nazionale di Geofisica e Vulcanologia, Roma, Italy*

² *Géoazur, CNRS, Observatoire de la Côte d'Azur, Université Côte d'Azur, Valbonne, France*

³ *Dipartimento di Fisica e Astronomia (DIFA), Alma Mater Studiorum Università di Bologna, Italy*

This is a pre-copyedited, author-produced PDF of an article accepted for publication in *Geophysical Journal International* following peer review. The version of record is available online at <https://doi.org/10.1093/gji/ggac263>.

SUMMARY

The computation of the Love numbers for a spherically symmetric self-gravitating viscoelastic Earth is a classical problem in global geodynamics. Here we revisit the problem of the numerical evaluation of loading and tidal Love numbers in the static limit for an incompressible planetary body, adopting a Laplace inversion scheme based upon the Post-Widder formula as an alternative to the traditional viscoelastic normal modes method. We also consider, within the same framework, complex-valued, frequency-dependent Love numbers that describe the response to a periodic forcing, which are paramount in the study of the tidal deformation of planets. Furthermore, we numerically obtain the time-derivatives of Love numbers, suitable for modeling geodetic signals in response to surface loads variations. A number of examples are shown, in which time and frequency-dependent Love numbers are evaluated for the Earth and planets adopting realistic rheological profiles. The numerical solution scheme is implemented in ALMA³ (the plAnetary Love nuMbers cALculator, version 3), an upgraded open-source Fortran 90 program that computes the Love numbers for radially layered planetary bodies with a wide range of rheologies, including transient laws like Andrade or Burgers.

Key words: Surface loading – Love numbers – Tides and planetary waves – Planetary interiors.

1 INTRODUCTION

Love numbers, first introduced by A.E.H. Love in 1911, provide a complete description of the response of a planetary body to external, surface or internal perturbations. In his seminal work, Love (1911) defined the Love numbers (LN) in the context of computing the radial deformation and the perturbation of gravity potential for an elastic, self-gravitating, homogeneous sphere that is subject to the gravitational pull of a tide-raising body. This definition has been subsequently extended by Shida (1912) to include also horizontal displacements. In order to describe the response to surface loads, an additional set of LNs, dubbed *loading Love numbers*, has been introduced in order to describe the Earth's response to surface loads (see *e.g.*, Munk & MacDonald, 1960; Farrell, 1972) and today they are routinely used in the context of the Post Glacial Rebound problem (Spada et al., 2011). In a similar way, *shear Love numbers* represent the response to a shear stress acting on the surface (Saito, 1978) while *dislocation Love numbers* describe deformations induced by internal point dislocations (see *e.g.*, Sun & Okubo, 1993).

The LN formalism has been originally defined in the realm of purely elastic deformations, for spherically symmetric Earth models consistent with global seismological observations. However, invoking the Correspondence Principle in linear viscoelasticity (see *e.g.*, Christensen, 1982), the LNs can be generalized to anelastic models in a straightforward way. Currently, viscoelastic LNs are a key ingredient of several geophysical applications involving the time-dependent response of a spherically symmetric Earth model to surface loads or endogenous perturbations. For example, they are essential to the solution of the Sea Level Equation (Farrell & Clark, 1976) and are exploited in current numerical implementations of the Glacial Isostatic Adjustment (GIA) problem, either on millennial (see *e.g.*, Spada & Melini, 2019) or on decadal time scale (see *e.g.*, Melini et al., 2015).

Since LNs depend on the internal structure of a planet and on its constitution, they can provide a means of establishing constraints on some physical parameters of the planet interior on the basis of geodetic measurements or astronomic observations (see *e.g.*, Zhang, 1992; Kellermann et al., 2018). For tidal periodic perturbations, complex LNs can be defined in the frequency domain, accounting for both the amplitude and phase lag of the response to a given tidal frequency (Williams & Boggs, 2015). Frequency-domain LNs are widely used to constrain the interior structure of planetary bodies on the basis of observations of tidal amplitude and phase lag (see *e.g.*, Sohl et al., 2003; Dumoulin et al., 2017; Tobie et al., 2019), to study the state of stress of satellites induced by tidal forcings (see *e.g.*, Wahr et al., 2009) or to investigate the tidal response of the giant planets (see *e.g.*, Gavrilov & Zharkov, 1977).

Viscoelastic LNs for a spherically symmetric, radially layered, self-gravitating planet are tradition-

ally computed within the framework of the “viscoelastic normal modes” method introduced by Peltier (1974), which relies upon the solution of Laplace-transformed equilibrium equations using the formalism of elastic propagators. As discussed by Spada & Boschi (2006) and Melini et al. (2008), this approach becomes progressively less feasible as the detail of the rheological model is increased or if complex constitutive laws are considered. Several workarounds have been proposed in the literature to avoid these shortcomings (see, e.g., Rundle, 1982; Friederich & Dalkolmo, 1995; Riva & Vermeersen, 2002; Tanaka et al., 2006). Among these, the Post-Widder Laplace inversion formula (Post, 1930; Widder, 1934), first applied by Spada & Boschi (2006) to the evaluation of viscoelastic LNs for the Earth, has the advantage of maintaining unaltered the formal structure of the viscoelastic normal modes and of allowing for a straightforward implementation of complex rheological laws. For periodic loads, alternative numerical integration schemes similar to those developed by Takeuchi & Saito (1972) for the elastic problem (Na & Baek, 2011; Wang et al., 2012) have been applied to the viscoelastic case by integrating Fourier-transformed solutions (Tobie et al., 2005, 2019).

In this work, we revisit the Post-Widder approach to the evaluation of LNs with the aim of extending it to more general planetary models, relaxing some of the assumptions originally made by Spada & Boschi (2006). In particular, we introduce a layered core in the Post-Widder formalism and obtain analytical expressions for the time derivatives of LNs, needed to model geodetic velocities in response to the variation of surface loads. In this respect, our approach is complementary to that of Padovan et al. (2018), who derived a semi-analytical solution for the fluid LNs using the propagator formalism. We implement our results in `iALMA` (the plAnetary Love nuMbers cAlculator, version 3), an open-source code which extends and generalizes the program originally released by Spada (2008). `iALMA` introduces a range of new capabilities, including the evaluation of frequency-domain LNs describing the response to periodic forcings, suitable for studying tidal dissipation in the Earth and planets.

This paper is organized as follows. In Section 2 we give a brief outline of the theory underlying the computation of viscoelastic LNs and of the application of the Post-Widder Laplace inversion formula. In Section 3 we discuss some general aspects of `iALMA`, leaving the technical details to a User Manual. In Section 4 we validate `iALMA` through some benchmarks between our numerical results and available reference solutions. In Section 5 we discuss some numerical examples before drawing our conclusions in Section 6.

2 MATHEMATICAL BACKGROUND

The details of the Post-Widder approach to numerical Laplace inversion have been extensively discussed in previous works (see Spada & Boschi, 2006; Spada, 2008; Melini et al., 2008). In what

follows, we only give a brief account of the Post-Widder Laplace inversion method for the sake of illustrating how the new features of ALMA have been implemented within its context.

2.1 Viscoelastic normal modes

Closed-form analytical expressions for the LNs exist only for a few extremely simplified planetary models. The first is the homogeneous, self-gravitating sphere, often referred to as the “Kelvin sphere” (Thomson, 1863). The second is the two-layer, incompressible, non self-gravitating model that has been solved analytically by Wu & Ni (1996). For more complex models, LNs shall be computed either through fully numerical integration of the equilibrium equations, or by invoking semi-analytical schemes. Among the latter, the viscoelastic normal modes method, introduced by Peltier (1974), relies upon the solution of the equilibrium equations in the Laplace-transformed domain. Invoking the Correspondence Principle (Christensen, 1982) the equilibrium equations can be cast in a formally elastic form by defining a complex rigidity (μ) that depends on the rheology adopted and is a function of the Laplace variable s .

Following Spada & Boschi (2006), at a given harmonic degree l , the Laplace-transformed equations can be solved with standard propagator methods, and their solution at the planet surface ($r = a$) can be written in vector form as

$$\mathbf{x}(s) = \mathbf{f}(s) \mathbf{P}_1(s) \mathbf{J} \mathbf{P}_2(s) \mathbf{J}^{-1} \mathbf{b}; \quad (1)$$

where the tilde denotes Laplace-transformed quantities, $\mathbf{v}(s) = (\tilde{u}; \tilde{v}; \tilde{w})^T$ contains the l -th degree harmonic coefficients of the vertical (and horizontal) components of the displacement field and the incremental potential, $\mathbf{f}(s)$ is the Laplace-transformed time-history of the forcing term, \mathbf{P}_1 and \mathbf{P}_2 are appropriate 6x6 projection operators, \mathbf{J} is a 6x3 array that accounts for the boundary conditions at the core interface, \mathbf{b} is a three-component vector expressing the surface boundary conditions (either of loading or of tidal type). In Eq. (1) \mathbf{J} is a 6x6 array that propagates the solution from the core radius $r = c$ to the planet surface ($r = a$), which has the form:

$$\mathbf{J}(s) = \prod_{k=N}^1 \mathbf{Y}_k(r_{k+1}; s) \mathbf{Y}_k^{-1}(r_k; s); \quad (2)$$

where N is the number of homogeneous layers outside the planet core, r_k is the radius of the interface between the $(k-1)$ -th and k -th layer, with $r_1 \leq \dots \leq r_N$, $r_1 = c$ and $r_{N+1} = a$. In Eq. (2), $\mathbf{Y}_k(r; s)$ is the fundamental matrix that contains the six linearly independent solutions of the equilibrium equations valid in the k -th layer, whose expressions are given analytically in Sabadini et al. (1982). When incompressibility is assumed, the matrix $\mathbf{Y}_k(r; s)$ depends upon the rheological constitutive law through the functional form of the complex rigidity (μ), which replaces the elastic rigidity

of the elastic propagator (Wu & Peltier, 1982). Table 1 lists expressions for some rheological laws. For a fluid inviscid (i.e., zero viscosity) core, the array in Eq. (1) is a 6 × 3 interface matrix whose components are explicitly given by Sabadini et al. (1982); conversely, for a solid core, responds to the 6 × 3 portion of the fundamental matrix for the core $\mathbf{Y}_e(c; s)$ that contains the three solutions behaving regularly for $\omega \neq 0$.

From the solution $\mathbf{x}(s)$ obtained in (1), the Laplace-transformed Love numbers are defined as:

$$h_n(s) = \frac{m}{a} u_n(s) \tag{3}$$

$$l_n(s) = \frac{m}{a} v_n(s) \tag{4}$$

$$k_n(s) = 1 - \frac{m}{ag} \dot{v}_n(s); \tag{5}$$

where we have made the dependence explicit, m is the mass of the planet and g is the unperturbed surface gravitational acceleration (Farrell, 1972; Wu & Peltier, 1982). Using Cauchy's residue theorem, for Maxwell or generalized Maxwell rheologies Eqs. (3-5) can be cast in the standard normal modes form, which for an impulsive load $f(t) = 1$ reads

$$L_n(s) = L_n^e + \sum_{k=1}^{N_M} \frac{L_n^k}{s - s_n^k}; \tag{6}$$

where $L_n(s)$ denotes any of the three LNs, L_n^e is the elastic component of the LNs, the limit for $s \rightarrow \infty$, L_n^k are the viscoelastic components (residues), s_n^k are the (real and negative) roots of the secular equation $\text{Det}(\mathbf{P}_2(s)) = 0$, and where N_M is the number of viscoelastic normal modes, each corresponding to one root of the secular equation (Spada & Boschi, 2006). However, such standard form is not always available, since for some particular rheologies the complex rigidity cannot be cast in the form of a rational fraction (this occurs, for example, for the Andrade's rheology, see Table 1). This is one of the motivations for adopting non-conventional Laplace inversion formulas like the one discussed in next section.

2.2 Love numbers in the time domain

To obtain the time-domain LNs, $h_n(t)$, $l_n(t)$ and $k_n(t)$, it is necessary to perform the inverse Laplace transform of Eqs. (3-5). Within the viscoelastic normal-mode approach, this is usually accomplished through an integration over a (modified) Bromwich path in the complex plane, by invoking the residue theorem. In this case, the inversion of Eq. (6) yields the time-domain Love numbers in the form:

$$L_n(t) = L_n^e(t) + H(t) \sum_{k=1}^{N_M} L_n^k e^{s_n^k t}; \tag{7}$$

where $\delta(t)$ is the Dirac delta and $H(t)$ is the Heaviside step function defined by Eq. (14) below, and

an impulsive time history is assumed ($\delta(t) = 1$). As discussed by Spada & Boschi (2006), the traditional scheme of the viscoelastic normal modes suffers from a few but significant shortcomings that, with models of increasing complexity, effectively hinders a reliable numerical inverse transformation. Indeed, the application of the residue theorem demands the identification of the poles of the Laplace-transformed solutions (see Eqs. 3-5), which are the roots of the secular polynomial equation whose algebraic degree increases with the number of rheologically distinct layers. In addition, its algebraic complexity may be unpractical to handle, particularly for constitutive laws characterized by many material parameters.

As shown by Spada & Boschi (2006) and Spada (2008), a possible way to circumvent these difficulties is to compute the inverse Laplace transform through the Post-Widder (PW) formula (Post, 1930; Widder, 1934). We note, however, that other viable possibilities exist, as the one recently discussed by Michel & Boy (2021), who have employed Fourier techniques to avoid some of the problems inherent in the Laplace transform method. While Fourier techniques may be more appropriate to take complex rheologies into account, and are clearly more relevant to address Love numbers at tidal frequencies, the motivation of our approach is to address in a unified framework the computation of LNs describing both tidal and surface loads. $\mathbb{F}(s) = \mathcal{L}\{F(t)\}$ is the Laplace transform of $F(t)$, the PW formula gives an asymptotic approximation of the inverse Laplace transform of $\mathbb{F}(s)$ as a function of the n -th derivatives of $\mathbb{F}(s)$ evaluated along the real positive axis:

$$F(t) = \lim_{n \rightarrow \infty} \frac{(1)^n}{n!} \frac{n^{n+1}}{t} \frac{d^n}{ds^n} \mathbb{F}(s) \Big|_{s=\frac{n}{t}} \quad (8)$$

In general, an analytical expression for the n -th derivative of $\mathbb{F}(s)$ required in Eq. (8) is not available. By employing a recursive discrete approximation of the derivative and rearranging the corresponding terms, Gaver (1966) has shown that an equivalent expression is

$$F(t) = \lim_{n \rightarrow \infty} \frac{n \ln 2}{t} \frac{2n!}{n} \sum_{j=0}^{n-1} \frac{x^n}{(1)^j} \frac{n!}{j} \mathbb{F} \left(\frac{(n+j) \ln 2}{t} \right) \quad ; \quad (9)$$

where the inverse transform $F(t)$ is expressed in terms of samples of the Laplace transform $\mathbb{F}(s)$ on the real positive axis of the complex plane. Since for a stably stratified incompressible planet all the singularities of $\mathbb{F}(s)$ (Eq. 1) are expected to be located along the real negative axis that ensures the long-term gravitational stability (Vermeersen & Mitrovica, 2000), Eq. (9) provides a strategy for evaluating the time-dependent LNs without the numerical complexities associated with the traditional contour integration. However, as discussed by Valabate (2004), the numerical convergence of (9) is logarithmically slow, and the oscillating terms can lead to catastrophic loss of numerical precision. Stehfest (1970) has shown that, for practical applications, the convergence of Eq. (9) can be accelerated by re-writing it in the form

$$F(t) = \lim_{M \rightarrow \infty} \frac{\ln 2}{t} \sum_{j=1}^M \frac{2^j}{j!} F_j \frac{j \ln 2}{t} ; \quad (10)$$

where M is the order of the Gaver sequence and where the constants are

$$k_{j,M} = \left(\frac{1}{2} \right)^{M+k} \frac{\min(M,k)}{j^{M+1}} \frac{M!}{j!} \frac{2^j}{j!} \frac{j!}{k!} ; \quad (11)$$

with $\text{oor}(x)$ being the greatest integer less or equal to x . Eq. (10) can be applied to (1) to obtain an M -th order approximation of the time-domain solution vector:

$$\underline{x}^{(M)}(t) = \frac{\ln 2}{t} \sum_{j=1}^M \frac{2^j}{j!} \underline{x}_j \frac{j \ln 2}{t} ; \quad (12)$$

from which the time-domain LNs can be readily obtained according to Eqs. (3-5).

Recalling that the Laplace transform $\underline{F}(s)$ and that of its time derivative $\underline{F}'(s)$ are related by $L(\underline{F}'(t)) = sL(\underline{F}(t)) - \underline{F}(0)$ and being $\underline{x}(t) = 0$ for $t < 0$, it is also possible to write an asymptotic approximation for the time derivative of the solution:

$$\underline{x}'^{(M)}(t) = \frac{\ln 2}{t} \sum_{j=1}^M \frac{2^j}{j!} \underline{x}_j \frac{j \ln 2}{t} ; \quad (13)$$

from which the time derivative of the LNs $\dot{s}_n(t)$, $\dot{l}_n(t)$ and $\dot{k}_n(t)$ can be obtained according to Eqs. (3-5). The numerical computation of the time-derivatives of the LNs according to Eq. (13) is one of the new features introduced in **ALMA**.

The time dependence of the solution vector obtained through Eqs. (12-13) is also determined by the time history of the forcing term (either of loading or tidal type), whose Laplace transform $\underline{f}(s)$ appears in Eq. (1). If the loading is instantaneously switched on at $t=0$, its time history is represented by the Heaviside (left-continuous) step function

$$H(t) = \begin{cases} 0; & t < 0 \\ 1; & t > 0; \end{cases} \quad (14)$$

whose Laplace transform is

$$\mathcal{H}(s) = L(H(t)) = \frac{1}{s} ; \quad (15)$$

Since any piece-wise constant function can be expressed as a linear combination of shifted Heaviside step functions (see, e.g. Spada & Melini, 2019), LNs obtained assuming the loading time history in Eq. (14) can be used to compute the response to arbitrary piece-wise constant loads. However, for some applications, it may be more convenient to represent the load time history as a piece-wise linear function. It is easy to show that any such function can be written as a linear combination of shifted elementary ramp functions of length t_r , of the type

$$R(t) = \begin{cases} 0; & t \leq 0 \\ \frac{t}{t_r}; & 0 < t \leq t_r \\ 1; & t > t_r; \end{cases} \quad (16)$$

whose Laplace transform is

$$\mathcal{R}(s) = L(R(t)) = \frac{1}{s} - \frac{1 - e^{-st_r}}{st_r} \quad (17)$$

Laplace-transformed LNs corresponding to a step-wise or ramp-wise forcing time history can be obtained by setting $f(s) = \mathcal{H}(s)$ or $f(s) = \mathcal{R}(s)$ in Eq. (1). The ramp-wise forcing function defined by Eq. (16) is one of the new features introduced in **ALMA**.

2.3 Frequency dependent Love numbers

In the context of planetary tidal deformation, it is important to determine the response to an external periodic tidal potential. The previous version of **ALMA** was limited to the case of an instantaneously applied forcing. For periodic potentials, the time dependence of the forcing term has the oscillating form $e^{i\omega t}$, where

$$\omega = \frac{2\pi}{T} \quad (18)$$

is the angular frequency of the forcing term, T is the period of the oscillation and $i = \sqrt{-1}$ is the imaginary unit. In the time domain, the solution vector can be cast in the form

$$x_i(t) = x_i(t) * e^{i\omega t} \quad (19)$$

where $x_i(t)$ is the time-domain response to an impulsive (delta) load and the asterisk indicates the time convolution. Since the impulsive load is a causal function ($\delta(t) = 0$ for $t < 0$) and Eq. (19) can be expressed as

$$x_i(t) = e^{i\omega t} \int_0^t x_i(t') e^{-i\omega t'} dt' = x_0(i\omega) e^{i\omega t} \quad (20)$$

where $x_0(i\omega)$ is the Laplace transform of $x_i(t)$ evaluated at $s = i\omega$. By setting $f(s) = L(\delta(t)) = 1$ and $s = i\omega$ in Eq. (1), we obtain

$$x_0(i\omega) = P_1(i\omega) J + P_2(i\omega) J^{-1} b \quad (21)$$

Hence, in analogy with Eqs. (3-5), the frequency-domain h_n(ω), $l_n(\omega)$ and $k_n(\omega)$ are defined as

$$h_n(\omega) = \frac{m}{a} u_n(\omega) \quad (22)$$

$$l_n(\omega) = \frac{m}{a} v_n(\omega) \quad (23)$$

$$k_n(\omega) = 1 - \frac{m}{ag} v_n(\omega); \quad (24)$$

where $u_n(\omega)$, $v_n(\omega)$ and $w_n(\omega)$ are the three components of vector $\mathbf{u}_n(\omega) = (u_n; v_n; w_n)^T$.

Since the frequency-domain LNs are complex numbers, in general a phase difference exists between the variation of the external periodic potential and the planet response, due to the energy dissipation within the planetary mantle. If $L_n(\omega)$ is any of the three frequency-dependent LNs, the corresponding time-domain LNs are:

$$L_n(t) = L_n(\omega) e^{i\omega t} = |L_n(\omega)| e^{i(\omega t - \phi)}; \quad (25)$$

where the phase lag ϕ is

$$\tan \phi = \frac{\text{Im}(L_n(\omega))}{\text{Re}(L_n(\omega))}; \quad (26)$$

and $\text{Re}(z)$ and $\text{Im}(z)$ denote the real and the imaginary parts of z , respectively. A vanishing phase lag ($\phi = 0$) is only expected for elastic planetary models (for $\text{Im}(L_n(\omega)) = 0$), for which no dissipation occurs. We remark that the evaluation of the frequency-dependent Love numbers (22-24) does not require the application of the Post-Widder method outlined in Section 2.2, since in this case no inverse transform is to be evaluated.

Tidal dissipation is phenomenologically expressed in terms of the quality factor Q (Kaula, 1964; Goldreich & Soter, 1966), which according to Efroimsky & Lainey (2007) and Clausen & Tilgner (2015) is related to the phase lag through

$$Q(\omega) = \frac{1}{\sin \phi} = \frac{|L_2(\omega)|}{\text{Im}(L_2(\omega))}; \quad (27)$$

thus implying $Q = 1$ in the case of no dissipation. Tidal dissipation is often measured in terms of the ratio

$$\frac{|k_2|}{Q} = |k_2| \sin \phi = \text{Im } k_2; \quad (28)$$

For terrestrial bodies, the quality factor usually lies in a range between 10 and 500 (Goldreich & Soter, 1966; Murray & Dermott, 2000). We remark that the quality factor is a phenomenological parameter used when the internal rheology is unknown; if LNs are computed by means of a viscoelastic model, it may be more convenient to consider the imaginary part, which is directly proportional to dissipation (Segatz et al., 1988).

3 AN OVERVIEW OF ALMA

Here we briefly outline how the solution scheme described in previous section is implemented in ALMA, leaving the technical details and practical considerations to the accompanying User Manual. ALMA evaluates, for any given harmonic degree, the time-domain LNs $\{h_n(t); l_n(t); k_n(t)\}$,

their time derivatives ($\dot{h}_n(t); \dot{l}_n(t); \dot{k}_n(t)$) and the frequency-domain LNs ($h_n(\omega); l_n(\omega); k_n(\omega)$), either corresponding to surface loading or to tidal boundary conditions. While the original version of the code was limited to time-domain LNs, the other two outputs represent new capabilities introduced by ALMA. The planetary model can include, in principle, any number of layers in addition to a central core. Each of the layers can be characterized by any of the rheological laws listed in Table 1, while the core can also have a fluid inviscid rheology. As we show in Section 5 below, numerical solutions obtained with ALMA are stable even with models including a large number of layers, providing a way to approximate rheologies whose parameters are varying continuously with radius.

Time-domain LNs are computed by evaluating numerically Eqs. (12) and (13), assuming a time history of the forcing that can be either a step function (Eq. 14) or an elementary ramp function (Eq. 16). In the latter case, the duration of the loading phase can be configured by the user. Since Eqs. (12) and (13) are singular for $\omega = 0$, ALMA can compute time-domain LNs only for $\omega > 0$. In the “elastic limit”, the LNs can be obtained either by sampling them at a time that is much smaller than the characteristic relaxation times of the model, or by configuring the Hooke's elastic rheology for all the layers in the model. In the second case, the LNs will follow the same time history of the forcing. As discussed in Section 2, the sums in Eqs. (12) and (13) contain oscillating terms that can lead to loss of precision due to catastrophic cancellation (Spada & Boschi, 2006). To avoid the consequent numerical degeneration of the LNs, ALMA performs all computations in arbitrary-precision floating point arithmetic, using the Fortran FMLIB library (Smith, 1991, 2003).

When running ALMA, the user shall configure both the number of significant digits used by the FMLIB library and the order M of the Gaver sequence in Eqs. (12) and (13). As discussed by Spada & Boschi (2006) and Spada (2008), higher values of D and M ensure a better numerical stability and accuracy of the results, but come at the cost of rapidly increasing computation time. All the examples discussed in the next section have been obtained with parameters $D=28$ and $M=8$. While these values ensure a good stability in relatively simple models, a special care shall be devoted to numerical convergence in case of models with a large number of layers and/or when computing LNs to high harmonic degrees; in that case, higher values of D and M may be needed to attain stable results.

Complex-valued LNs are obtained by ALMA by directly sampling Eq. (21) at the requested frequencies ω , and therefore no numerical Laplace anti-transform is performed. While for frequency-domain LNs the numerical instabilities associated with the Post-Widder formula are avoided, the use of high-precision arithmetic may still be appropriate, especially in case of models including a large number of layers. ALMA does not directly compute the tidal phase lag, the quality factor Q nor the $k_2=Q$ ratio, which can be readily obtained from tabulated output values of the real and imaginary parts of LNs through Eqs. (26-28).

Although ALMA is still limited to spherically symmetric and elastically incompressible models, with respect to the version originally released by Spada (2008) now the program includes some new significant features aimed at increasing its versatility. These include: i) the evaluation of frequency-dependent loading and tidal Love numbers in response to periodic forcing; ii) the possibility of dealing with a layered core that includes fluid and solid portions; iii) the introduction of a ramp-shaped forcing function to facilitate the implementation of loading histories varying in a linear piecewise manner; iv) the implementation of the Andrade transient viscoelastic rheology often employed in the study of planetary deformations; v) the explicit evaluation of the derivatives of the Love numbers in the time domain to facilitate the computation of geodetic variations in deglaciated areas; vi) a short but exhaustive User Guide, and vii) a facilitated computation of frequency-dependent loading and tidal planetary Love numbers, with pre-defined and easily customizable rheological profiles for some terrestrial planets and moons.

4 BENCHMARKING ALMA

In the following we discuss a suite of numerical benchmarks for Love numbers computed by ALMA. First, we consider a uniform, incompressible, self-gravitating sphere with Maxwell rheology (the so-called “Kelvin sphere”) and compare tidal LNs computed numerically by ALMA with well known analytical results. Then, we test numerical results from ALMA by reproducing the viscoelastic LNs for an incompressible Earth model computed within the benchmark exercise by Spada et al. (2011). Finally, we discuss the impact of the incompressibility approximation assumption by comparing elastic and viscoelastic LNs for a realistic Earth model with recent numerical results by Michel & Boy (2021), who employ a compressible model.

4.1 The viscoelastic Kelvin sphere

Simplified planetary models for which closed-form expressions for the LNs are available are of particular relevance here, since they allow an analytical benchmarking of the numerical solutions discussed in Section 2 and provided by ALMA.

In what follows, we consider a spherical, homogeneous, self-gravitating model, often referred to as the “Kelvin sphere” (Thomson, 1863), which can be extended to a viscoelastic rheology in a straightforward manner. For example, adopting the complex modulus appropriate for the Maxwell rhe-

ology (see Table 1), for a Kelvin sphere of radius a , density ρ and surface gravity g , in the Laplace domain the harmonic degree $n = 2$ LNs take the form

$$\mathcal{L}_2(s) = \frac{L_f}{1 + \frac{2s}{s+1}}; \quad (29)$$

where L_2 stands for any of $(h_2; l_2; k_2)$, L_f is the “tidal limit” of $\mathcal{L}_2(s)$ (i.e., the value attained for $s \rightarrow 0$), the Maxwell relaxation time is

$$\tau = \frac{1}{\rho g a} \quad (30)$$

and

$$\tau^2 = \frac{19}{2} \frac{a^3}{g} \quad (31)$$

is a positive non-dimensional constant. Note that τ is a function of a and ρ , since for the homogeneous sphere $\rho = \frac{4}{3} G a$, where G is the universal gravitational constant.

After some algebra, (29) can be cast in the form

$$\mathcal{L}_2(s) = \frac{L_f}{1 + \frac{2s}{s+1}} = \frac{L_f}{1 + \frac{1-s}{s+1}}; \quad (32)$$

where for a tidal forcing, the tidal limits for degree $n = 2$ are $h_f = \frac{5}{2}$, $l_f = \frac{3}{4}$ and $k_f = \frac{3}{2}$ (see e.g., Lambeck, 1988) and where we have defined

$$\tau^0 = (1 + \frac{2}{s+1})^{-1} \quad (33)$$

From Eq. (32), the LNs in the time domain can be immediately computed analytically through an inverse Laplace transformation:

$$L_2(t) = \frac{L_f}{1 + \frac{2}{s+1}} (t) + H(t) \left[\frac{1}{s} - \frac{1}{s+1} \right] e^{-t/\tau^0}; \quad (34)$$

while for an external forcing characterized by a step-wise time-history, the $L_2^{(H)}(t)$ are obtained by a time convolution with the Heaviside function:

$$L_2^{(H)}(t) = L_2(t) * H(t); \quad (35)$$

that yields

$$L_2^{(H)}(t) = \frac{L_f}{1 + \frac{2}{s+1}} \left[1 + \frac{2}{s+1} \right] e^{-t/\tau^0}; \quad t > 0; \quad (36)$$

from which the time derivative $\frac{dL_2^{(H)}}{dt}(t)$ is readily obtained:

$$\frac{dL_2^{(H)}}{dt}(t) = \frac{L_f}{1 + \frac{2}{s+1}} \left[-\frac{1}{\tau^0} \right] e^{-t/\tau^0}; \quad t > 0; \quad (37)$$

F1

In Figure 1a, the dotted curves show the (blue) and the k_2 (red) tidal LN of harmonic degree $n = 2$ obtained by a configuration of ALMA that reproduces the Kelvin sphere (the parameters are given in the Figure caption). The LNs, shown as a function of time, are characterized by two

asymptotes corresponding to the elastic and fluid limits, respectively, and by a smooth transition in between. The solid curves, obtained by the analytical expression given by Eq. (36), show an excellent agreement with the ALMA^{A} numerical solutions. The same holds for the time-derivatives of these LNs, considered in Figure 1b, where the analytical LNs (solid lines) are computed according to Eq. (37).

The frequency response of the Kelvin sphere for a periodic tidal potential can be obtained by settings $\nu = i\omega$ in Eq. (29), which after rearranging gives:

$$L_2(\omega) = \frac{L_f}{1 + \frac{\omega^2}{\omega_0^2}} + i \frac{2\omega}{1 + \frac{\omega^2}{\omega_0^2}} \quad (38)$$

which remarkably depends upon ω and ω_0 only through the ω/ω_0 product. Therefore, a change in the relaxation time τ shall result in a shift of the frequency response of the Kelvin sphere, leaving its shape unaltered.

Using Eq. (38) in (26), the phase lag turns out to be:

$$\tan \delta = \frac{2\omega}{1 + \frac{\omega^2}{\omega_0^2}} \quad (39)$$

where it is easy to show that for frequency

$$\omega_0 = \frac{1}{\rho \tau} \quad (40)$$

the maximum phase lag δ_{\max} is attained, with

$$\tan \delta_{\max} = \frac{2}{1 + \frac{\omega_0^2}{\omega_0^2}} \quad (41)$$

By using Eq. (38) into (27), for the Kelvin sphere the quality factor is

$$Q_K(\omega) = \frac{1}{1 + \frac{\omega^2}{\omega_0^2} + \frac{1}{4} \frac{\omega^4}{\omega_0^4}} \quad (42)$$

which at $\omega = \omega_0$ attains its minimum value

$$Q_{\min} = 1 + \frac{2}{2} \quad (43)$$

In Figure 2a, the dotted curve shows the phase lag δ as a function of the tidal period $T = 2\pi/\omega$, F_2 obtained by the same configuration ALMA^{A} described in the caption of Figure 1. The solid line corresponds to the analytical expression (39) which can be obtained from Eq. (39), showing once again an excellent agreement with the numerical results (dotted). Figure 2b compares numerical results obtained from ALMA^{A} for Q with the analytical expression for $Q_K(T)$ obtained from (42). By using in Eq. (40) the numerical values of ρ and τ assumed in Figures 1 and 2, the period $T_0 = 2\pi/\omega_0$ is found to scale with viscosity as

$$T_0 = (3.06 \text{ kyr}) \frac{1}{10^{21} \text{ Pa s}} \quad (44)$$

so that for $\tau = 10^{21} \text{ Pa s}$, representative of the Earth's mantle bulk viscosity (see Mitrovića,

1996; Turcotte & Schubert, 2014), the maximum phase lag $\tau_0 = 41:9$ and the minimum quality factor $Q_{\min} = 1:5$ are attained for $T_0 = 3$ kyr, consistent with the results shown in Figure 2.

4.2 Community-agreed Love numbers for an incompressible Earth model

Due to the relevance of viscoelastic Love numbers in a wide range of applications in Earth science, several numerical approaches for their evaluation have been independently developed and proposed in literature. This ignited the interest on benchmark exercises, in which a set of agreed numerical results can be obtained and different approaches and methods can be cross-validated. Here we consider a benchmark effort that has taken place in the framework of the Glacio-Isostatic Adjustment community (Spada et al., 2011), in which a set of reference viscoelastic Love numbers for an incompressible, spherically symmetric Earth model has been derived through different numerical approaches, including viscoelastic normal modes, spectral- finite elements and finite elements. This allows us to validate our numerical results by implementing the M3-L70-V01 Earth model described in Table of Spada et al. (2011), which includes a fluid inviscid core, three mantle layers with Maxwell viscoelastic rheology and an elastic lithosphere, and comparing the set of Love numbers with reference results from the benchmark exercise.

F3 Figure 3 shows elastic ($h_n^{(e)}$, $l_n^{(e)}$, $k_n^{(e)}$) and fluid Love numbers ($h_n^{(f)}$, $l_n^{(f)}$, $k_n^{(f)}$), both for the loading and tidal cases, computed by ALMA for the M3-L70-V01 Earth model in the range of harmonic degrees $2 \leq n \leq 250$. The elastic and fluid limits have been simulated by ALMA by sampling the time-dependent Love numbers at $t_e = 10^{-5}$ kyr and $t_f = 10^{10}$ kyr, respectively. Reference results from Spada et al. (2011), represented by solid lines in Figure 3, are practically indistinguishable from results obtained with ALMA over the whole range of harmonic degrees, demonstrating the reliability of the numerical approach employed.

F4 Figure 4 shows time-dependent Love numbers ($h_n(t)$, $l_n(t)$ and $k_n(t)$), for both the loading and tidal cases, computed by ALMA for harmonic degrees $2 \leq n \leq 5$ and for t between 10^{-3} and 10^5 kyr, a time range that encompasses the complete transition between the elastic and fluid limits. Also in this case, numerical results obtained by ALMA (shown by symbols) are coincident with the reference Love numbers from Spada et al. (2011), represented by solid lines.

4.3 Viscoelastic Love numbers for a PREM-layered Earth model

In this last benchmark, we compare numerical results from ALMA with reference viscoelastic Love numbers for a realistic Earth model which accounts for an elastically compressible rheology, in order to assess its

importance when modeling the tidal and loading response of a large planetary body. In the context of Earth rotation, the role of compressibility has been addressed by Vermeersen et al. (1996); the reader is also referred to Sabadini et al. (2016) for a broader presentation of the problem and to Renaud & Henning (2018) for a discussion of the effects of compressibility in the realm of planetary modelling.

Here we focus on numerical results recently obtained by Michel & Boy (2021), who employed Fourier techniques to compute frequency-dependent viscoelastic LNs for periodic forcings both of loading and tidal types. They have adopted an Earth model with the elastic structure of PREM (Preliminary Reference Earth Model, Dziewonski & Anderson, 1981) and a fully liquid core, and replaced the outer oceanic layer with a solid crust layer, adjusting crustal density in such a way to keep the total Earth mass unchanged. Following Michel & Boy (2021), we have built a discretised realization of PREM suitable for ALMA with a liquid core and 28 homogeneous mantle layers, which has been used to obtain the numerical results discussed below.

Figure 5 compares elastic Love numbers obtained by Michel & Boy (2021) in the range of harmonic degrees between $n = 2$ and $n = 10,000$ with those computed with ALMA. The largest difference between the two sets of LNs can be seen for the loading case (Figure 5a), where the assumption of incompressibility leads to a significant underestimation of deformation across the whole range of harmonic degrees. Incompressible elasticity leads to an underestimation also of the loading LN (Figure 5b), although the differences are much smaller and limited to the lowest harmonic degrees. Conversely, for the tidal response (Figures 5c and 5d) the two sets of LNs turn out to be almost overlapping, suggesting a minor impact of elastic compressibility on tidal deformations.

In Figure 6 we consider a periodic load and compare viscoelastic tidal LNs computed with ALMA with corresponding results from Michel & Boy (2021). Consistently with the elastic case, we see that the incompressibility approximation used in ALMA generally results in smaller modeled deformations across the whole range of forcing periods. The largest differences are found on $|k_2|$ (6a) and reach the 20% level in the range of periods between 10^5 and 10^6 days, while on $|k_2|$ (6b) the differences are much smaller, reaching the 10% level in the same range of periods. Similarly, for the phase lags (Figures 6c and 6d) we find a larger difference for k_2 , with the phase lag being remarkably insensitive to compressibility up to forcing periods of the order of 10^5 days.

5 EXAMPLES OF ALMA APPLICATIONS

In this Section we consider four applications showing the potential of ALMA in different contexts. First, we will discuss the k_2 tidal Love number of Venus, based upon a realistic layering for the

interior of this planet. Second, we shall evaluate the tidal LNs for a simple model of the Saturn's moon Enceladus, in order to show how an internal fluid layer can be simulated as a low-viscosity Newtonian fluid rheology and how a depth-dependent viscosity in a conductive shell may be approximated using a sequence of thin homogeneous layers. Third, we will evaluate a set of loading LNs suitable for describing the transient response of the Earth to the melting of large continental ice sheets. As a last example, we will demonstrate how ALMA can simulate the tidal dissipation on the Moon using two recent interior models based on seismological data. While these numerical experiments are put in the context of state-of-the-art planetary interior modeling, we remark that they are aimed only at illustrating the modeling capabilities of ALMA.

5.1 Tidal deformation of Venus

The planet Venus is often referred to as “Earth's twin planet”, since its size and density differ only by 5% from those of the Earth. These similarities lead to the expectation that the chemical composition of the Earth and Venus may be similar, with an iron-rich core, a magnesium silicate mantle and a silicate crust (Kovach & Anderson, 1965; Lewis, 1972; Anderson, 1980). Despite these similarities, there is a lack of constraints on the internal structure of Venus. Therefore, its density and rigidity profiles are often assumed to be a re-scaled version of the Preliminary Reference Earth Model (PREM) of Dziewonski & Anderson (1981), accounting for the difference in the planet's radius and mass, as in Aitta (2012). One of the main observational constraints on the planet's interior, along its mass and moment of inertia, is its J_2 tidal LN. The current observational estimate for Venus is 0.295 ± 0.066 (2 σ formal), and it has been inferred from Magellan and Pioneer Venus orbiter spacecraft data (Konopliv & Yoder, 1996). However, due to uncertainties, it is not possible to discriminate between a liquid and a solid core (Dumoulin et al., 2017).

Here we use ALMA to reproduce results obtained by means of the Venus model referred to as T_5^{hot} by Dumoulin et al. (2017), based on the “hot temperature profile” from Armann & Tackley (2012), having a composition and hydrostatic pressure from the PREM model of Dziewonski & Anderson (1981). The viscosity of the mantle of Venus is fixed and homogeneous; the crust is elastic ($\nu = 1$), the core is assumed to be inviscid ($\nu = 0$) and the rheology of the mantle follows Andrade's law (see Table 1). The parameters of the model have been volume-averaged into the core, the lower mantle, the upper mantle and the crust. The calculation is performed at the tidal period of 58.4 days (Cottureau et al., 2011). In the work of Dumoulin et al. (2017) is computed by integrating the radial functions associated with the gravitational potential, as defined by Takeuchi & Saito (1972), hence the simplified formulation of Saito (1974) relying on the radial function is employed.

The method is derived from the classical theory of elastic body deformation and the energy density integrals commonly used in the seismological community. One of the main differences between their computation and the results presented here is the assumption about compressibility, since Dumoulin et al. (2017) use a compressible planetary model, while ALMA an incompressible rheology is always assumed. In Figure 7, the two curves show the tidal LN corresponding to Andrade creep F7 parameters $n = 0.2$ and $n = 0.3$ as a function of mantle viscosity for the tidal period of 58.4 days. Each of the vertical red segments corresponds to the interval values obtained by Dumoulin et al. (2017) for discrete mantle viscosity values $\eta = 10^{19}$, 10^{20} , 10^{21} and 10^{22} Pas, respectively, and for a range of the Andrade creep parameter in the interval between 0.2 and 0.3. The grey shaded area illustrates the most recent observed values of k_2 according to Konopliv & Yoder (1996) to an uncertainty of 2 formal σ . Figure 7 shows that the values obtained with ALMA for the T_5^{hot} Venus model fit well with the lower boundary of the compared study for each of the discrete mantle viscosity values if an Andrade creep parameter $n = 0.3$ is assumed, while for $n = 0.2$ the modeled k_2 slightly exceeds the upper boundary of Dumoulin et al. (2017).

5.2 The tidal response of Enceladus

The scientific interest on Enceladus has gained considerable momentum after the 2005 Cassini flybys, which confirmed the icy nature of its surface and evidenced the existence of water-rich plumes emerging from the southern polar regions (Porco et al., 2006; Ivins et al., 2020). These hint to the existence of a subsurface ocean, heated by tidal dissipation in the core, where physical conditions allowing life could be possible, in principle (for a review, see Hemingway et al., 2018). The interior structure of Enceladus has been thoroughly investigated in literature on the basis of observations of its gravity field (Iess et al., 2014), tidal deformation and physical librations (see, e.g., Cadek et al., 2016), setting constraints on the possible structure of the ice shell and of the underlying liquid ocean (Roberts & Nimmo, 2008), and on the composition of its core (Roberts, 2015). Lateral variations in the crustal thickness of Enceladus have been inferred in studies about the isostatic response of the satellite using gravity and topography data as constraints (Cadek et al., 2016; Beuthe et al., 2016; Cadek et al., 2019) and in works dealing with the computation of deformation and dissipation (see, e.g., Cadek et al., 2016, 2019; Beuthe, 2018, 2019). Indeed, from all the above studies, it clearly emerges that a full insight into the tidal dynamics of Enceladus could be only gained adopting 3D models of its internal structure.

While a thorough investigation of the signature of the interior structure of Enceladus on its tidal response is far beyond the scope of this work, here we set up a simple spherically symmetric model

with the purpose of illustrating how the LNs for a planetary body including a fluid internal layer like Enceladus can be computed with ALM \tilde{A} , and how a radially-dependent viscosity structure can be approximated with homogeneous layers. We define a spherically symmetric model including an homogeneous inner solid core of radius 192 km (Hemingway et al., 2018), surrounded by a liquid water layer and an outer icy shell, and investigate the sensitivity of the tidal LNs to the thickness of the ice layer, along the lines of Roberts & Nimmo (2008) and Beuthe (2018). In our setup, the core is modeled as a homogeneous elastic body with rigidity $\mu = 4 \cdot 10^{10}$ Pa and whose density is adjusted to ensure that, when varying the thickness of the ice shell, the average bulk density of the model is kept constant at $\rho_b = 1610$ kg m $^{-3}$. Since in ALM \tilde{A} a fluid inviscid rheology can be prescribed only for the core, we approximate the ocean layer as a low viscosity Newtonian fluid (10^4 Pas). The ice shell is modeled as a conductive Maxwell body whose viscosity profile depends on the temperature T according to the Arrhenius law:

$$\eta(T) = \eta_m \exp \left(\frac{E_a}{R_g T} \right) \quad (45)$$

where E_a is the activation energy, R_g is the gas constant, T_m is the temperature at the base of the ice shell and η_m is the ice viscosity at $T = T_m$. Following Beuthe (2018), we use $E_a = 59.4$ J=(mol K), $\eta_m = 10^{13}$ Pas and $T_m = 273$ K, and assume that the temperature inside the ice shell varies with radius r according to

$$T(r) = T_m \frac{r - a}{r_b - a} \frac{r_b - r}{r_b - a}; \quad (46)$$

where r_b is the bottom radius of the ice shell and $T_s = 59$ K is the average surface temperature. Since in ALM \tilde{A} the rheological parameters must be constant inside each layer, we discretize the radial viscosity profile given by Eq. (45) using a onion-like structure of homogeneous spherical shells. To assess the sensitivity of results to the choice of discretization resolution, we perform three numerical experiments in which the thickness of ice layers is set to 0.25, 0.5 and 1 km. The ice and water densities are set to $\rho_i = 930$ kg m $^{-3}$ and $\rho_w = 1020$ kg m $^{-3}$, respectively, while the ice rigidity is set to $\mu_i = 3.5 \cdot 10^9$ Pa, a value consistent with evidence from tidal flexure of marine ice (Vaughan, 1995) and laboratory experiments (Cole & Durell, 1995).

Figure 8a shows the elastic tidal LNs, l_2 and k_2 for the Enceladus model discussed above as a function of the thickness of the ice shell. The elastic tidal response is strongly dependent on the ice thickness, with the k_2 LNs decreasing from 0.090 for a 10 km-thick shell to 0.015 for a 50 km-thick shell. It is of interest to compare these results with elastic LNs obtained by Beuthe (2018) in the uniform-shell approximation. It turns out that the LN shown in Figure 8a is slightly smaller than corresponding results from Beuthe (2018), with relative differences between 5-10% level, consistently with their estimate of the effect of incompressibility. Figure 8b shows the real and

imaginary parts of the h_2 tidal LN as a function of the thickness of the ice layer for a periodic load of period $T = 1.37$ days, which corresponds to the shortest librational oscillation of Enceladus (Rambaux et al., 2010). As discussed above, for this numerical experiment we implemented a radially-variable viscosity profile by discretizing Eq. (45) into a series of uniform layers. Solid and dashed lines in Figure 8b show results obtained with a discretization step of 0.5 km and 1.0 km, respectively; we verified that with a step of 0.25 km the results are virtually identical to those obtained with a step of 0.5 km. The effect of the discretization is evident only on the imaginary part, where a coarse layer size of 1 km leads to a significant overestimation of $\text{Im}(h_2)$ if the ice shell is thinner than 15 km. By a visual comparison of the results of Figure 8b with Figure 4 of Beuthe (2018), we can see that the imaginary part of h_2 is well reproduced, while the real part is underestimated by the same level we found for the elastic LNs; this difference is likely to be attributed to the incompressibility approximation adopted in ALMA.

5.3 Loading Love numbers for transient rheologies in the Earth's mantle

Loading Love numbers are key components in models of the response of the Earth to the spatio-temporal variation of surface loads, including the ongoing deformation due to the melting of the late Pleistocene ice complexes (see, e.g., Peltier & Drummond, 2008; Purcell et al., 2016), the present-day and future response to climate-driven melting of ice sheets and glaciers (Bamber & Riva, 2010; Slangen, 2012), and deformations induced by the variation of hydrological loads (Bevis et al., 2016; Silverii et al., 2016). Evidence from Global Navigation Satellite System measurements of the time-dependent surface deformation point to a possible transient nature of the mantle in response to the regional-scale melting of ice sheets and to large earthquakes (see Pollitz, 2003, 2005; Nield et al., 2014; Qiu et al., 2018). Here, it is therefore of interest to present the outcomes of some numerical experiments in which ALMA is configured to compute the time-dependent loading Love Number assuming a transient rheology in the mantle. Numerical estimates of $h_n(t)$ and of its time derivative $\dot{h}_n(t)$ would be needed, for instance, to model the response to the thickness variation of a disc-shaped surface load, as discussed by Bevis et al. (2016).

In Figure 9 we show the time evolution of the $h_2(t)$ loading LN for $n = 2, 10$ and 100 , comparing the response obtained assuming the VM5a viscosity model of Peltier & Drummond (2008), which is fully based on a Maxwell rheology, with those expected if VM5a is modified introducing a transient rheology in the upper mantle layers. An Heaviside time history for the load is adopted throughout. In model VM5a-BG we assumed a Burgers bi-viscous rheological law in the upper mantle with τ_1 and $\tau_2 = \tau_1 = 0.1$ (see Table 1), while in model VM5a-AD an Andrade rheology (Cottrell, 1996)

with creep parameter $n = 0.3$ has been assumed for the upper mantle. For 2 (Figure 9a) the responses obtained with the three models almost overlap. Indeed, for long wavelengths (by Jean's rule, the wavelength corresponding to harmonic degree $n = \frac{2a}{\lambda}$, where a is Earth's radius) the response to surface loads is mostly sensitive to the structure of the lower mantle, where the three variants of VM5a considered here have the same rheological properties. Conversely (Figure 9b) we see a slightly faster response to the loading for both transient models in the time range between 0.01 and 1 kyr. For $n = 100$, the transient response of VM5a-BG and VM5a-AD becomes even more enhanced between 0.01 and 10 kyr. It is worth to note that, for times less than 10 kyr, the two transient versions of VM5a almost yield identical responses, suggesting that an Andrade rheology in the Earth's upper mantle might explain the observed vertical transient deformations in the same way as a Burgers rheology. The differences between the three models are more evident in Figure 10, where we use ALMA for computing the time derivatives $\dot{s}_n(t)$ (this option was not available in previous versions of the program). Compared with the Maxwell model, the transient ones show a significantly larger initial rate of vertical displacement, that differ significantly for Burgers and Andrade. The three rheologies provide comparable responses only 0.1 kyrs after loading. We shall remark, however, that the incompressibility approximation employed in ALMA has a significant impact on the Love Number, as we discussed in Section 4.3, so the results shown above must be taken with caution, and a more detailed analysis of the impact of compressibility on the time evolution of LNs would be in order.

F10

5.4 Tidal dissipation on the Moon

The Moon is the extraterrestrial body for which the most detailed information about the internal structure is available. In addition to physical constraints from observations of tidal deformation (Williams et al., 2014), seismic experiments deployed during the Apollo missions (Nunn et al., 2020) provided instrumental recordings of moonquakes which allowed the formulation of a set of progressively refined interior models (see, e.g. Heffels et al., 2021).

In this last numerical experiment, we configured ALMA to compute tidal LNs for the Moon according to the two interior models proposed by Weber et al. (2011, W11 hereafter) and Garcia et al. (2011, 2012, G12 hereafter). Profiles of density and rigidity for models W11 and G12 are shown in Figure 11, with the most notable difference being that the former assumes an inner solid core and a fluid outer core, while the latter contains an undifferentiated fluid core. We emphasize that model G12 includes 70 rheological layers in the mantle and crust, demonstrating the stability of ALMA with densely-layered planetary models. For both models, we assumed a Maxwell rheology in the crust and

F11

the mantle, with a viscosity of 10^{20} Pas. A more realistic approach has been followed by Nimmo et al. (2012), who have modelled the Moon's Love numbers and dissipation adopting an extended Burgers model for the mantle, which also accounts for transient tidal deformations (Faul & Jackson, 2015). Such rheological model is not incorporated in the current release of ALMA, but it can be implemented by the user modifying the source code in order to compute the corresponding complex rigidity modulus $\mu^*(s)$. The fluid core has been modeled as a Newtonian fluid with viscosity 10^4 Pas while in the inner core, for model W11, we used a Maxwell rheology with a viscosity of 10^{18} Pas, a value within the estimated ranges for the viscosity of the Earth inner core (Buffett, 1997; Dumberry & Mound, 2010; Koot & Dumberry, 2011). Following the lines of Harada et al. (2014, 2016) and Organowski & Dumberry (2020), we defined a 50 km thick low-viscosity zone (LVZ) at the base of the mantle and computed the k_2 tidal LNs as a function of the LVZ viscosity for a forcing period $P = 27.212$ days.

For both W11 and G12 models, Figure 12 shows the dependence on the LVZ viscosity of the F_{12} k_2 tidal LN (Figure 12a), of its phase lag angle (12b) and of the quality factor Q (12c). With the considered setup, for a LVZ viscosity smaller than 10^{15} Pas the tidal response of the two models is almost coincident, while for higher viscosities model G12 predicts a stronger tidal dissipation. Shaded gray areas in frames (12a) and (12c) show dependence intervals for experimental estimates of k_2 (Williams et al., 2014) and Q (Williams & Boggs, 2015). With both models we obtain values of Q within the 1- interval for an LVZ viscosity smaller than about 10^{15} Pas; interestingly, for that LVZ viscosity the G12 model predicts a quality factor within the measured range, while model W11 would require a slightly higher LVZ viscosity (10^{16} Pas). Of course, a detailed assessment of the ability of the two models to reproduce the observed tidal LNs would be well beyond the scope of this work, and several additional parameters potentially affecting the tidal response (e.g. the LVZ thickness or the core radius) would need to be considered.

6 CONCLUSIONS

We have revisited the Post-Widder approach in the context of evaluating viscoelastic Love numbers and their time derivatives for arbitrary planetary models. Our results are the basis of a new version of ALMA, a user friendly Fortran program that computes the Love numbers of a multi-layered, self-gravitating, spherically symmetric, incompressible planetary model characterized by a linear viscoelastic rheology. ALMA can be suitably employed to solve a wide range of problems, either involving the surface loading or the tidal response of a rheologically layered planet. By taking advantage of the Post-Widder Laplace inversion method, the evaluation of the time-domain Love numbers is simplified, avoiding some of the limitations of the traditional viscoelastic normal mode approach.

Differently from previous implementations (Spada, 2008), `ALMA` can evaluate both time-domain and frequency-domain Love numbers, for an extended set of linear viscoelastic constitutive equations that also include a transient response, like Burgers or Andrade rheologies. Generalized linear rheologies that until now have been utilized in at geometry like the one characterizing the extended Burgers model (Ivins et al., 2020) could be possibly implemented as well modifying the source code, if the corresponding analytical expression of the complex rigidity modulus is available. Furthermore, `ALMA` can compute the time-derivatives of the Love numbers, and can deal with step-like and ramp-shaped forcing functions. The resulting Love numbers can be linearly superposed to obtain the planet response to arbitrarily time evolving loads. Numerical results from `ALMA` have been benchmarked with analytical expressions for a uniform sphere and with a reference set of viscoelastic LNs for an incompressible Earth model (Spada et al., 2011). The well-known limitations of the incompressibility approximation in modeling deformations of large terrestrial bodies have been quantitatively assessed by a comparison between numerical outputs `ALMA` and viscoelastic LNs recently obtained by Michel & Boy (2021) for a realistic, compressible Earth model. The versatility of `ALMA` has then been demonstrated by a few examples, in which the Love numbers and some associated quantities like the quality factor have been evaluated for some multi-layered models of planetary interiors characterized by complex rheological profiles and by densely-layered internal structures.

ACKNOWLEDGMENTS

We thank the Associate Editor Gael Choblet and two anonymous Reviewers for their very constructive comments that considerably helped improving our manuscript. We have benefited from discussion with all scientists involved in the project “LDLR – Lunar tidal Deformation from Earth-based and orbital Laser Ranging”, funded by the French ANR and the German agency DGF. We are indebted to Steve Vance, Saikiran Tharimena, Marshall Styczinski and Bruce Bills for encouragement and advice. DM is funded by a INGV (Istituto Nazionale di Geofisica e Vulcanologia) 2020-23 “ricerca libera” research grant and partly supported by the INGV project Pianeta Dinamico 2021-22 Tema 4 KINDLE (grant no. CUP D53J19000170001), funded by the Italian Ministry of University and Research “Fondo nazionale al rilancio degli investimenti delle amministrazioni centrali dello Stato e allo sviluppo del Paese, Legge 145/2018”. CS is funded by a PhD grant of the French Ministry of Research and Innovation. CS also acknowledges the ANR, project number ANR-19-CE31-0026 project LDLR (Lunar tidal Deformation from Earth-based and orbital Laser Ranging). GS is funded by a FFABR (Finanziamento delle Attività Base di Ricerca) grant of MIUR (Ministero dell'Istruzione, dell'Università e della Ricerca).

della Ricerca) and by a RFO research grant of DIFA (Dipartimento di Fisica e Astronomia “Augusto Righi”) of the Alma Mater Studiorum Università di Bologna.

Data availability. Source code of the ALMA version used to obtain numerical results presented in this work is available as a supplementary material. The latest version ALMA can be downloaded from <https://github.com/danielemelini/ALMA3>. The data underlying plots shown in this article are available upon request to the corresponding author.

References

- Aitta, A., 2012. Venus' internal structure, temperature and core composition, *Journal of Geophysics*, 218(2), 967–974.
- Anderson, D. L., 1980. Tectonics and composition of Venus, *Geophysical Research Letters*, 7(1), 101–102.
- Armann, M. & Tackley, P. J., 2012. Simulating the thermochemical magmatic and tectonic evolution of Venus's mantle and lithosphere: Two-dimensional models, *Geophys. Res.*, 117, E12003
- Bamber, J. & Riva, R., 2010. The sea level fingerprint of recent ice mass loss, *The Cryosphere*, 4(4), 621–627.
- Beuthe, M., 2018. Enceladus's crust as a non-uniform thin shell: I tidal deformations, *Journal of Geophysics*, 302, 145–174.
- Beuthe, M., 2019. Enceladus's crust as a non-uniform thin shell: II tidal dissipation, *Journal of Geophysics*, 332, 66–91.
- Beuthe, M., Rivoldini, A., & Trinh, A., 2016. Enceladus's and Dione's coating ice shells supported by minimum stress isostasy, *Geophysical Research Letters*, 43(19), 10–088.
- Bevis, M., Melini, D., & Spada, G., 2016. On computing the geoelectric response to a disk load, *Geophysical Journal International*, 205(1), 1804–1812.
- Buffett, B. A., 1997. Geodynamic estimates of the viscosity of the Earth's inner core, *Nature*, 388, 1476–4687.
- Cadek, O., Tobie, G., Van Hoolst, T., Mészáros, M., Choblet, G., Lefèvre, A., Mitri, G., Baland, R.-M., Behouk, M., Bourgeois, O., & Trinh, A., 2016. Enceladus's internal ocean and ice shell constrained from Cassini gravity, shape, and libration data, *Geophysical Research Letters*, 43(11), 5653–5660.
- Cadek, O., Sotik, O., Behouk, M., Choblet, G., Tobie, G., & Hron, J., 2019. Long-term stability of Enceladus' uneven ice shell, *Journal of Geophysics*, 319, 476–484.
- Christensen, R., 1982. *Theory of Viscoelasticity*, Dover, Mineola, New York.

- Clausen, N. & Tilgner, A., 2015. Dissipation in rocky planets for strong tidal forcing, *Astronomy & Astrophysics*, 584, A60.
- Cole, D. M. & Durell, G. D., 1995. The cyclic loading of saline ice, *Philosophical Magazine A* 72(1), 209–229.
- Cottureau, L., Rambaux, N., Lebonnois, S., & Souchay, J., 2011. The various contributions in Venus rotation rate and LOD, *Astronomy & Astrophysics*, 531, A45.
- Cottrell, A. H., 1996. Andrade creep, *Philosophical Magazine Letters*, 73(1), 35–36.
- Dumberry, M. & Mound, J., 2010. Inner core–mantle gravitational locking and the super-rotation of the inner core, *Geophysical Journal International*, 181(2), 806–817.
- Dumoulin, C., Tobie, G., Verhoeven, O., Rosenblatt, P., & Rambaux, N., 2017. Tidal constraints on the interior of Venus, *Journal of Geophysical Research: Planets*, 122(6), 1338–1352.
- Dziewonski, A. M. & Anderson, D. L., 1981. Preliminary reference Earth model, *Physics of the Earth and Planetary Interiors*, 25(4), 297–356.
- Efroimsky, M. & Lainey, V., 2007. Physics of bodily tides in terrestrial planets and the appropriate scales of dynamical evolution, *Journal of Geophysical Research: Planets*, 112, E12003.
- Farrell, W., 1972. Deformation of the Earth by surface loads, *Reviews of Geophysics*, 10(3), 761–797.
- Farrell, W. & Clark, J., 1976. On postglacial sea level, *Geophysical Journal International*, 46, 647–667.
- Faul, U. & Jackson, I., 2015. Transient creep and strain energy dissipation: An experimental perspective, *Annual Review of Earth and Planetary Sciences*, 43, 541–569.
- Friederich, W. & Dalkolmo, J., 1995. Complete synthetic seismograms for a spherically symmetric earth by a numerical computation of the Green's function in the frequency domain, *Geophysical Journal International*, 122(2), 537–550.
- Garcia, R. F., Gagnepain-Beyneix, J., Chevrot, S., & Lognonné, P., 2011. Very preliminary reference Moon model, *Physics of the Earth and Planetary Interiors*, 188(1), 96–113.
- Garcia, R. F., Gagnepain-Beyneix, J., Chevrot, S., & Lognonné, P., 2012. Erratum to “Very Preliminary Reference Moon Model”, by R.F. Garcia, J. Gagnepain-Beyneix, S. Chevrot, P. Lognonné [Phys. Earth Planet. Inter. 188 (2011) 96–113], *Physics of the Earth and Planetary Interiors*, 202–203, 89–91.
- Gaver, D. P., 1966. Observing stochastic processes, and approximate transform in operations, *Research*, 14(3), 444–459.
- Gavrilov, S. & Zharkov, V., 1977. Love numbers of the giant planets, *Icarus*, 32(4), 443–449.
- Goldreich, P. & Soter, S., 1966. Q in the solar system, *Icarus*, 5(1), 375–389.
- Harada, Y., Goossens, S., Matsumoto, K., Yan, J., Ping, J., Noda, H., & Haruyama, J., 2014. Strong

- tidal heating in an ultralow-viscosity zone at the core–mantle boundary of the Moon, *Nature Geoscience* **7**, 569–572.
- Harada, Y., Goossens, S., Matsumoto, K., Yan, J., Ping, J., Noda, H., & Haruyama, J., 2016. The deep lunar interior with a low-viscosity zone: Revised constraints from recent geodetic parameters on the tidal response of the Moon, *Earth and Planetary Science Letters* **276**, 96–101.
- Heffels, A., Knapmeyer, M., Oberst, J., & Haase, I., 2021. Re-evaluation of Apollo 17 Lunar Seismic Profiling Experiment data including new LROC-derived coordinates for explosive packages 1 and 7, at Taurus-Littrow, Moon, *Planetary and Space Science* **206**, 105307.
- Hemingway, D., Less, L., Tajeddine, R., & G., G. T., 2018. The interior of Enceladus, Callisto and the Icy Moons of Saturn, pp. 57–77, eds Schenk, P. M., Clark, R. N., Howett, C. J. A., Verbiscer, A. J., & Waite, J. H., University of Arizona, Tucson.
- Less, L., Stevenson, D., Parisi, M., Hemingway, D., Jacobson, R., Lunine, J., Nimmo, F., Armstrong, J., Asmar, S., Ducci, M., et al., 2014. The gravity field and interior structure of Enceladus, *Science* **344**(6179), 78–80.
- Ivins, E. R., Caron, L., Adhikari, S., Larour, E., & Scheinert, M., 2020. A linear viscoelasticity for decadal to centennial time scale mantle deformation, *Reports on Progress in Physics* **83**(10), 106801.
- Kaula, W. M., 1964. Tidal dissipation by solid friction and the resulting orbital evolution, *Reviews of Geophysics* **2**(4), 661–685.
- Kellermann, C., Becker, A., & Redmer, R., 2018. Interior structure models and fluid Love numbers of exoplanets in the super-Earth regime, *Astronomy & Astrophysics* **615**, A39.
- Konopliv, A. & Yoder, C., 1996. Venusian k_2 tidal Love number from Magellan and PVO tracking data, *Geophysical Research Letters* **23**(14), 1857–1860.
- Koot, L. & Dumberry, M., 2011. Viscosity of the Earth's inner core: Constraints from nutation observations, *Earth and Planetary Science Letters* **308**(3), 343–349.
- Kovach, R. L. & Anderson, D. L., 1965. The interiors of the terrestrial planets, *Journal of Geophysical Research* **70**(12), 2873–2882.
- Lambeck, K., 1988. The earth's variable rotation: some geophysical causes, *Symposium—International Astronomical Union*, vol. 128, pp. 1–20, Cambridge University Press.
- Lewis, J. S., 1972. Metal/silicate fractionation in the solar system, *Earth and Planetary Science Letters* **15**(3), 286–290.
- Love, A. E. H., 1911. *Some Problems of Geodynamics: Being an Essay to which the Adams Prize in the University of Cambridge was Adjudged in 1909*, CUP Archive.
- Melini, D., Cannelli, V., Piersanti, A., & Spada, G., 2008. Post-seismic rebound of a spherical

- Earth: new insights from the application of the Post-Widder inversion formula, *Geophysical Journal International* 174(2), 672–695.
- Melini, D., Gegout, P., King, M., Marzeion, B., & Spada, G., 2015. On the rebound: Modeling Earth's ever-changing shape, *EOS* 96(15), 14–17.
- Michel, A. & Boy, J.-P., 2021. Viscoelastic Love numbers and long-period geophysical effects, *Geophysical Journal International* 228(2), 1191–1212.
- Mitrovica, J. X., 1996. Haskell [1935] revisited, *Journal of Geophysical Research: Solid Earth* 101(B1), 555–569.
- Munk, W. H. & MacDonald, G. J., 1960. *The Rotation of the Earth: A Geophysical Discussion* Cambridge Univ. Press, New York.
- Murray, C. D. & Dermott, S. F., 2000. *Solar System Dynamics* Cambridge University Press.
- Na, S.-H. & Baek, J., 2011. Computation of the Load Love number and the Load Green's function for an elastic and spherically symmetric earth, *Journal of the Korean Physical Society* 58(5), 1195–1205.
- Nield, G., Barletta, V., Bordoni, A., King, M., Whitehouse, P., Clarke, P., Domack, E., Scambos, T., & Berthier, E., 2014. Rapid bedrock uplift in the Antarctic Peninsula explained by viscoelastic response to recent ice unloading, *Earth and Planetary Science Letters* 247, 32–41.
- Nimmo, F., Faul, U., & Garnero, E., 2012. Dissipation at tidal and seismic frequencies in a melt-free Moon, *Journal of Geophysical Research: Planets* 117(E9).
- Nunn, C., Garcia, R. F., Nakamura, Y., Marusiak, A. G., Kawamura, T., Sun, D., Margerin, L., Weber, R., Drilleau, M., Wiczorek, M. A., Khan, A., Rivoldini, A., Lognonné, P., & Zhu, P., 2020. Lunar seismology: A data and instrumentation review, *Space Science Reviews* 216(89).
- Organowski, O. & Dumberry, M., 2020. Viscoelastic relaxation within the Moon and the phase lead of its Cassini state, *Journal of Geophysical Research: Planets* 125(7), e2020JE006386.
- Padovan, S., Spohn, T., Baumeister, P., Tosi, N., Breuer, D., Csizmadia, S., Hellard, H., & Sohl, F., 2018. Matrix-propagator approach to compute tidal Love numbers and applicability to extrasolar planets, *Astronomy & Astrophysics* 620, A178.
- Peltier, W. & Drummond, R., 2008. Rheological stratification of the lithosphere: A direct inference based upon the geodetically observed pattern of the glacial isostatic adjustment of the North American continent, *Geophysical Research Letters* 35(16).
- Peltier, W. R., 1974. The impulse response of a Maxwell Earth, *Review of Geophysics and Space Physics* 12(4), 649–669.
- Pollitz, F. F., 2003. Transient rheology of the uppermost mantle beneath the Mojave Desert, California, *Earth and Planetary Science Letters* 215(1-2), 89–104.

- Pollitz, F. F., 2005. Transient rheology of the upper mantle beneath central Alaska inferred from the crustal velocity field following the 2002 Denali earthquake. *Journal of Geophysical Research: Solid Earth*, 110(B8).
- Porco, C. C., Helfenstein, P., Thomas, P., Ingersoll, A., Wisdom, J., West, R., Neukum, G., Denk, T., Wagner, R., Roatsch, T., et al., 2006. Cassini observes the active south pole of Enceladus. *Science*, 311(5766), 1393–1401.
- Post, E. L., 1930. Generalized differentiation. *Transactions of the American Mathematical Society* 32(4), 723–781.
- Purcell, A., Tregoning, P., & Dehecq, A., 2016. An assessment of the ICE6GM5a glacial isostatic adjustment model. *Journal of Geophysical Research: Solid Earth* 121(5), 3939–3950.
- Qiu, Q., Moore, J. D. P., Barbot, S., Feng, L., & Hill, E. M., 2018. Transient rheology of the Sumatran mantle wedge revealed by a decade of great earthquakes. *Nature Communications*, 9(1), 995.
- Rambaux, N., Castillo-Rogez, J. C., Williams, J. G., & Karatekin, Ö., 2010. Librational response of Enceladus. *Geophysical Research Letters* 37(4).
- Renaud, J. P. & Henning, W. G., 2018. Increased tidal dissipation using advanced rheological models: Implications for Io and tidally active exoplanets. *The Astrophysical Journal* 857(2), 98.
- Riva, R. E. M. & Vermeersen, L. L. A., 2002. Approximation method for high-degree harmonics in normal mode modelling. *Geophysical Journal International* 151(1), 309–313.
- Roberts, J. H., 2015. The fluffy core of Enceladus. *Nature*, 528, 54–66.
- Roberts, J. H. & Nimmo, F., 2008. Tidal heating and the long-term stability of a subsurface ocean on Enceladus. *Nature*, 454(7182), 675–689.
- Rundle, J. B., 1982. Viscoelastic-gravitational deformation by a rectangular thrust fault in a layered Earth. *Journal of Geophysical Research: Solid Earth* 87(B9), 7787–7796.
- Sabadini, R., Yuen, D. A., & Boschi, E., 1982. Polar wandering and the forced responses of a rotating, multilayered, viscoelastic planet. *Journal of Geophysical Research: Solid Earth* 87(B4), 2885–2903.
- Sabadini, R., Vermeersen, B., & Cambiotti, G., 2006. *Global dynamics of the Earth*. Springer.
- Saito, M., 1974. Some problems of static deformation of the Earth. *Journal of Physics of the Earth* 22(1), 123–140.
- Saito, M., 1978. Relationship between tidal and load Love numbers. *Journal of Physics of the Earth* 26(1), 13–16.
- Segatz, M., Spohn, T., Ross, M., & Schubert, G., 1988. Tidal dissipation, surface heat flow, and structure of viscoelastic models of Io. *Nature*, 334(6137), 187–206.
- Shida, T., 1912. *On the elasticity of the Earth and the Earth's crust*. Kyoto Imperial University.

- Silverii, F., D'Agostino, N., Métois, M., Fiorillo, F., & Ventafridda, G., 2016. Transient deformation of karst aquifers due to seasonal and multiyear groundwater variations observed by GPS in southern Apennines (Italy), *Journal of Geophysical Research: Solid Earth* **121**(11), 8315–8337.
- Slangen, A., 2012. Modelling regional sea-level changes in recent past and future. PhD. thesis, Utrecht University, the Netherlands.
- Smith, D., 2003. Using multiple-precision arithmetic. *Computing in Science Engineering* **5**(4), 88–93.
- Smith, D. M., 1991. Algorithm 693: A FORTRAN Package for Floating-Point Multiple-Precision Arithmetic, *ACM Trans. Math. Software* **17**(2), 273–283.
- Sohl, F., Hussmann, H., Schwentker, B., Spohn, T., & Lorenz, R. D., 2003. Interior structure models and tidal Love numbers of Titan. *Journal of Geophysical Research: Planets* **108**(E12).
- Soucek, O., Hron, J., Bhounkova, M., & Cadek, O., 2016. Effect of the tiger stripes on the deformation of Saturn's moon Enceladus. *Geophysical Research Letters* **43**(14), 7417–7423.
- Soucek, O., Bhounkova, M., Cadek, O., Hron, J., Tobie, G., & Choblet, G., 2019. Tidal dissipation in Enceladus' uneven, fractured ice shell. *Nature* **568**, 218–231.
- Spada, G., 2008. ALMA, a Fortran program for computing the viscoelastic Love numbers of a spherically symmetric planet. *Computers & Geosciences* **34**(6), 667–687.
- Spada, G. & Boschi, L., 2006. Using the Post-Widder formula to compute the Earth's viscoelastic Love numbers. *Geophysical Journal International* **166**(1), 309–321.
- Spada, G. & Melini, D., 2019. SELEN (SELEN version 4.0): a Fortran program for solving the gravitationally and topographically self-consistent sea-level equation in glacial isostatic adjustment modeling. *Geoscientific Model Development* **12**(12), 5055–5075.
- Spada, G., Barletta, V. R., Klemann, V., Riva, R., Martinec, Z., Gasperini, P., Lund, B., Wolf, D., Vermeersen, L., & King, M., 2011. A benchmark study for glacial isostatic adjustment codes. *Geophysical Journal International* **185**(1), 106–132.
- Stehfest, H., 1970. Algorithm 368: Numerical inversion of Laplace transforms. *Communications of the ACM* **13**(1), 47–49.
- Sun, W. & Okubo, S., 1993. Surface potential and gravity changes due to internal dislocations in a spherical earth—I. Theory for a point dislocation. *Geophysical Journal International* **114**(3), 569–592.
- Takeuchi, H. & Saito, M., 1972. Seismic surface wave methods in computational physics. *Methods in computational physics* **1**, 217–295.
- Tanaka, Y., Okuno, J., & Okubo, S., 2006. A new method for the computation of global viscoelastic post-seismic deformation in a realistic earth model (I)—vertical displacement and gravity variation,

- Geophysical Journal International, **164**(2), 273–289.
- Thomson, W., 1863. XXVII. On the rigidity of the earth. *Philosophical Transactions of the Royal Society of London*, **153**, 573–582.
- Tobie, G., Mocquet, A., & Sotin, C., 2005. Tidal dissipation within large icy satellites: Applications to Europa and Titan. *Earth and Planetary Science Letters*, **230**(1–2), 534–549.
- Tobie, G., Grasset, O., Dumoulin, C., & Mocquet, A., 2019. Tidal response of rocky and ice-rich exoplanets. *Astronomy & Astrophysics*, **630**, A70.
- Turcotte, D. L. & Schubert, G., 2014. *Geodynamics - Applications of Continuum Physics to Geological Problems*. Cambridge University Press.
- Valkó, P. P. & Abate, J., 2004. Comparison of sequence accelerators for the Gaver method of numerical Laplace transform inversion. *Computers & Mathematics with Applications*, **48**(3), 629–636.
- Vaughan, D. G., 1995. Tidal exure at ice shelf margins. *Journal of Geophysical Research: Solid Earth*, **100**(B4), 6213–6224.
- Vermeersen, L. & Mitrovica, J., 2000. Gravitational stability of spherical self-gravitating relaxation models. *Geophysical Journal International*, **142**(2), 351–360.
- Vermeersen, L. L. A., Sabadini, R., & Spada, G., 1996. Compressible rotational deformation. *Geophysical Journal International*, **126**, 735–761.
- Wahr, J., Selvens, Z. A., Mullen, M. E., Barr, A. C., Collins, G. C., Selvens, M. M., & Pappalardo, R. T., 2009. Modeling stresses on satellites due to nonsynchronous rotation and orbital eccentricity using gravitational potential theory. *Earth and Planetary Science Letters*, **280**(1), 188–206.
- Wang, H., Xiang, L., Jia, L., Jiang, L., Wang, Z., Hu, B., & Gao, P., 2012. Load Love numbers and Green's functions for elastic Earth models PREM, iasp91, ak135, and modified models with refined crustal structure from Crust 2.0. *Computers & Geosciences*, **49**, 190–199.
- Weber, R. C., Lin, P.-Y., Garnero, E. J., Williams, Q., & Lognonné, P., 2011. Seismic detection of the lunar core. *Science*, **331**(6015), 309–312.
- Widder, D. V., 1934. The inversion of the Laplace integral and the related moment problem. *Transactions of the American Mathematical Society*, **39**(1), 107–200.
- Williams, J. G. & Boggs, D. H., 2015. Tides on the Moon: Theory and determination of dissipation. *Journal of Geophysical Research: Planets*, **120**(4), 689–724.
- Williams, J. G., Konopliv, A. S., Boggs, D. H., Park, R. S., Yuan, D.-N., Lemoine, F. G., Goossens, S., Mazarico, E., Nimmo, F., Weber, R. C., Asmar, S. W., Melosh, H. J., Neumann, G. A., Phillips, R. J., Smith, D. E., Solomon, S. C., Watkins, M. M., Wieczorek, M. A., Andrews-Hanna, J. C., Head, J. W., Kiefer, W. S., Matsuyama, I., McGovern, P. J., Taylor, G. J., & Zuber, M. T., 2014. Lunar interior properties from the GRAIL mission. *Journal of Geophysical Research: Planets*, **119**(7), 1207–1224.

1546–1578.

Wu, P. & Ni, Z., 1996. Some analytical solutions for the viscoelastic gravitational relaxation of a two-layer non-self-gravitating incompressible spherical earth. *Geophysical Journal International* 126(2), 413–436.

Wu, P. & Peltier, W., 1982. Viscous gravitational relaxation. *Geophysical Journal International* 70(2), 435–485.

Zhang, C., 1992. Love numbers of the Moon and of the terrestrial planets. *Earth, Moon, and Planets* 56(3), 193–207.

Table 1. Complex rigidities $\hat{\sigma}(s)$ for the linear viscoelastic rheologies implemented in ALMA. Here, μ is the elastic rigidity, η is the Newtonian viscosity, μ_2 and η_2 are the rigidity and viscosity of the transient element in the bi-viscous Burgers rheology, respectively. In the Andrade rheological law, τ is the creep parameter while $\Gamma(x)$ is the Gamma function.

Rheological law	Complex rigidity $\hat{\sigma}(s)$
Hooke	μ
Maxwell	$\frac{\mu s}{s + \frac{\mu}{\eta}}$
Newton	ηs
Kelvin	$\frac{\mu \eta}{\mu + \eta s}$
Burgers	$\frac{\mu s + \frac{\mu_2 \eta_2}{s}}{s^2 + s \left(\frac{\mu}{\eta} + \frac{\mu_2}{\eta_2} \right) + \frac{\mu_2 \eta_2}{s}}$
Andrade	$\frac{1}{1 + \frac{1}{s} + \left(\frac{\mu}{\eta} + 1 \right) \frac{1}{s} + \frac{\tau}{s}}$

Figure 1. (a) Comparison between numerical (dotted) and analytical solutions (solid) for the k_2 tidal LNs of a Kelvin sphere with Maxwell rheology having radius = 6371 km, density = $5.514 \cdot 10^3 \text{ kg m}^{-3}$, rigidity = $1.46 \cdot 10^{11} \text{ Pa}$ and viscosity = 10^{21} Pas . (b) The same, for the time derivatives of the LNs. Note that the time axis is logarithmic.

Figure 2. Comparison between numerical (dotted) and analytical solutions (solid) for the tidal phase lag and quality factor Q (b) for the $n = 2$ tidal LNs of a Kelvin sphere with Maxwell rheology, using the same parameters detailed in the caption of Figure 1.

Figure 3. Elastic (red) and fluid (blue) Love numbers as a function of the harmonic degree for the Earth model M3-L70-V01 defined in Spada et al. (2011). Top (a-c) and bottom frames (d-f) show Love numbers for loading and tidal forcing, respectively. Symbols show numerical results obtained with ~~ALMA~~ while solid lines represent reference results from the benchmark exercise by Spada et al. (2011).

Figure 4. Time-dependent viscoelastic Love numbers for the M3-L70-V01 Earth model at long spatial wavelength (harmonic degrees $n = 5$). Top panels (frames a-c) and bottom ones (d-f) show Love numbers for a loading and tidal forcing, respectively. The time history of the load is an Heaviside unit step function. Symbols show numerical results obtained with `WLM` while solid lines represent reference results from the benchmark exercise by Spada et al. (2011).

f

Figure 5. Comparison between elastic Love numbers (left panels) and k_n (right) obtained by Michel & Boy (2021) with numerical results from ALMA. In both cases, the Earth model has the elastic structure of PREM in the crust and in the mantle, while the core is modeled as an uniform, inviscid fluid. Top and bottom panels show loading and tidal Love numbers, respectively.

Figure 6. Comparison between the modulus of tidal Love numbers h_2 and k_2 (top panels) and corresponding phase lags (bottom panels) obtained by Michel & Boy (2021) for a periodic forcing with numerical results from ALMA. The Earth model has the elastic structure of PREM and a Maxwell rheology with viscosity 10^{21} Pas is assumed in the mantle.

Figure 7. Tidal Love Number k_2 as a function of the mantle viscosity η_M for the internal model T_5^{hot} of Dumoulin et al. (2017). The two curves correspond to numerical results assuming Andrade creep parameters $n = 0.2$ and $n = 0.3$, respectively. Red vertical segments represent the range of the estimates obtained by Dumoulin et al. (2017), while the grey shaded area represents the most recent observed value of k_2 and its uncertainty according to Konopliv & Yoder (1996).

Figure 8. Elastic harmonic degree 2 tidal Love numbers for Enceladus (a) as a function of the thickness of the ice shell. In (b), real and imaginary parts of the viscoelastic tidal Love number for a forcing period of 1.73 days are shown. Solid lines and dashed lines correspond to discretization steps for the ice shell of 0.50 and 1.00 km, respectively. Please note that k_2 has been multiplied by a factor 10 to improve readability.




Figure 9. Loading Love number $h_n(t)$ for $n = 2$ (frame a), $n = 10$ (b) and $n = 100$ (c), obtained with the VM5a viscosity model by Peltier & Drummond (2008) and with two variants that assume Burgers (VM5a-BG) or Andrade (VM5a-AD) rheologies in the upper mantle layers.




Figure 10. Time-derivative of the loading Love number $\dot{h}_n(t)$ for harmonic degrees $n = 2, 10$ and 100 , adopting the rheological models described in the caption of Figure 9.




Figure 11. Radial profiles of density (a) and rigidity (b) for the Moon models by Weber et al. (2011) (W11, blue) and Garcia et al. (2011, 2012) (G12, red). Models W11 and G12 include 10 and 71 homogeneous layers, respectively.




Figure 12. Modulus of the tidal Love number $|k_2|$ for the Moon (frame a), phase lag (b) and quality factor (c) as a function of the LVZ viscosity, for a forcing period $T = 27.212$ days. Blue and red curves correspond to the Moon models by Weber et al. (2011) and Garcia et al. (2011, 2012) shown in Figure 11. Shaded areas in frames (a) and (c) correspond to the 1- σ confidence intervals for measured values of k_2 and Q according to Williams & Boggs (2015).