

The Surface faulting produced by the 30 October 2016 Mw 6.5 Central Italy earthquake: the Open EMERGEIO Working Group experience

Daniela Pantosti and the Open EMERGEIO Working Group Team
Istituto Nazionale di Geofisica e Vulcanologia, Italy

The October 30, 2016 (06:40 UTC) Mw 6.5 earthquake occurred about 28 km NW of Amatrice village as the result of upper crust normal faulting on a nearly 30 km-long, NW-SE oriented, SW dipping fault system in the Central Apennines. This earthquake is the strongest Italian seismic event since the 1980 Mw 6.9 Irpinia earthquake. The Mw 6.5 event was the largest shock of a seismic sequence, which began on August 24 with a Mw 6.0 earthquake and also included a Mw 5.9 earthquake on October 26, about 9 and 35 km NW of Amatrice village, respectively. Field surveys of coseismic geological effects at the surface started within hours of the mainshock and were carried out by several national and international teams of earth scientists (about 120 people) from different research institutions and universities coordinated by the EMERGEIO Working Group of the Istituto Nazionale di Geofisica e Vulcanologia. This collaborative effort was focused on the detailed recognition and mapping of: 1) the total extent of the October 30 coseismic surface ruptures, 2) their geometric and kinematic characteristics, 3) the coseismic displacement distribution along the activated fault system, including subsidiary and antithetic ruptures.

The huge amount of collected data (more than 8000 observation points of several types of coseismic effects at the surface) were stored, managed and shared using a specifically designed spreadsheet to populate a georeferenced database. More comprehensive mapping of the details and extent of surface rupture was facilitated by Structure-from-Motion photogrammetry surveys by means of several helicopter flights.

An almost continuous alignment of ruptures about 30 km long, N150/160 striking, mainly SW side down was observed along the already known active Mt. Vettore - Mt. Bove fault system. The mapped ruptures occasionally overlapped those of the August 24 Mw 6.0 and October 26 Mw 5.9 shocks. The coincidence between the observed surface ruptures and the trace of active normal faults mapped in the available geological literature is noteworthy. The field data collected suggest a complex coseismic surface faulting pattern along closely-spaced, parallel or subparallel, overlapping or step-like synthetic and antithetic fault splays. The cumulative surface faulting length has been estimated in about 40 km. The maximum vertical offset is significant, locally exceeding 2 meters along the Mt. Vettore Fault, measured both along bedrock fault planes and free-faces affecting unconsolidated deposits. This enormous collaborative experience has a twofold relevance, on the one side allowed to document in high detail the earthquake ruptures before Winter would destroy them, on the other represent the first large European experience for coseismic effects survey that we should use a leading case to establish a coseismic effects European team to get ready to respond to future seismic crises at the European level.

The Open EMERGEO Working Group Team

Pucci Stefano, De Martini Paolo Marco, Civico Riccardo, Cinti Francesca Romana, Pizzimenti Luca, Nappi Rosa, Ricci Tullio, Villani Fabio, Caciagli Marco, Di Naccio Deborah, Brunori Carlo Alberto, Sapia Vincenzo, Fracassi Umberto, Cucci Luigi, Nave Rosella, Alessio Giuliana, Sciarra Alessandra, Gaudiosi Germana, Alfonsi Laura, Vallone Roberto, Tarabusi Gabriele, Tarquini Simone, Del Carlo Paola, Mazzarini Francesco, Smedile Alessandra, Azzaro Raffaele, Moro Marco, Amoroso Sara, Kastelic Vanja, Carafa Michele, Tiberti Mara Monica, Branca Stefano, Burrato Pierfrancesco, Bertagnini Antonella, Braun Thomas, De Ritis Riccardo, Gori Stefano, Vannoli Paola, Falcucci Emanuela, Pagliuca Nicola, Lisi Arianna, Venuti Alessandra, Mariucci Maria Teresa, Buttinelli Mauro, Bisson Marina, Patera Antonio, Montone Paola, Pezzo Giuseppe, Pinzi Stefania, Pignone Maurizio, Pantosti Daniela (**Istituto Nazionale di Geofisica e Vulcanologia, INGV, Italy**);
Eutizio Vittori, Luca Guerrieri, Anna Maria Blumetti, Marco Amanti, Fabrizio Galluzzo, Claudio Campobasso, Pio Di Manna, Valerio Comerci (**Istituto Superiore per la Protezione e la Ricerca Ambientale, ISPRA, Italy**);
Michetti Alessandro Maria, Livio Franz, Ferrario Maria Francesca, Frigerio Chiara, Silvia Terrana, Andrea Pozzi, Gilberto Binda, Roberto Gambillara, Livio Bonadeo (**Università dell'Insubria, Italy**);
Luigi Piccardi (**Consiglio Nazionale delle Ricerche, Institute of Geosciences and Earth Resources (IGG), Firenze, Italy**);
Emanuele Tondi, Domenico Aringoli, Chiara Invernizzi, Farabollini Piero, Marco Materazzi, Pietro Paolo Pierantoni, Silvia Paggi, Gilberto Pambianchi, Eugenio Pistolesi, Titiano Volatili, Miller Zambrano, Danica Jablonska, Deiana Giovanni (**Università di Camerino, Italy**);
Fabrizio Agosta (**Università della Basilicata, Italy**);
Luca Falconi, Salomon Hailemikael, Marco Proposito, Claudio Puglisi (**Agenzia nazionale per le nuove tecnologie, l'energia e lo sviluppo economico sostenibile, ENEA, Italy**);
Michele Saroli, Lorenzo Lo Sardo (**Università degli Studi di Cassino e del Lazio Meridionale, DICeM, Italy**);
Giusy Lavecchia, Francesco Brozzetti, Paolo Boncio, Daniele Cirillo, Federica Ferrarini, Francesca Liberi, Eugenio Auciello, Simone Bello (**Università Chieti-Pescara, Centro Interuniversitario per l'Analisi Sismotettonica Tridimensionale, CRUST-UR Chieti, Italy**);
Bruno Pace, Alessandro Valentini (**Università Chieti-Pescara, DiSPUTer Chieti, Italy**);
Gerald Roberts, Iezzi Francesco (**Birkbeck University of London, United Kingdom**);
Laura Gregory, Huw Goodall, Luke Wedmore (**University of Leeds, United Kingdom**);
Joanna Faure Walker, Mildon Zoe (**University College London, United Kingdom**);
Ken McCaffrey (**Durham University, United Kingdom**);
Max Wilkinson (**Geospatial Research Ltd, United Kingdom**);
Stéphane Baize, Hervé Jomard, Oona Scooti (**Institut de Radioprotection et Sûreté Nucléaire ; BERSSIN ; Fontenay-aux-Roses, France**);
Lucilla Benedetti, Jules Fleury, Magali Rizza, Jim Tesson, Eugénie Pérouse (**Aix-Marseille Université, CEREGE CNRS-IRD UMR 34, Aix en Provence, France**);
Nathalie Feuillet, Eric Jacques, Yann Klinger, Arthur Delorme (**Institut de Physique du Globe de Paris, Sorbonne Paris Cité, Paris, France**);
Matthieu Ferry (**Géosciences Montpellier, CNRS, Montpellier, France**);
Frédérique Leclerc (**Université Côte d'Azur, CNRS, Observatoire de la Côte d'Azur, IRD, Géoazur**);
Jérôme van der Woerd, Jean-Philippe Malet, Marc Odin, Julien Point (**Université de Strasbourg, CNRS, Institut de Physique du Globe de Strasbourg UMR 7516, Strasbourg, France**);
Pierrick Bornemann, Skupinski Grzegorz (**Université de Strasbourg, CNRS, Laboratoire Image Ville Environnement UMR 7362, Strasbourg, France**);
Luca Del Rio (**Università degli Studi di Roma "La Sapienza", Italy**);
Irene Puliti (**Università di Perugia, Italy**);
Licia Ripamonti, Marzio Roncoroni (**SOGIN, Italy**)