



Shallow subsurface imaging of the Piano di Pezza active normal fault (central Italy) using high-resolution refraction and electrical resistivity tomography coupled with time-domain electromagnetic data

Villani, F. (1), Tulliani, V. (2), Sapia, V. (3), Fierro, E. (4), Civico, R. (3), Baccheschi, P. (1), Di Giulio, G. (1), Vassallo, M. (1), Pantosti, D. (3)

- (1) Istituto Nazionale di Geofisica e Vulcanologia, via dell'Arcivescovado 8, 67100 L'Aquila, Italy. Email: fabio.villani@ingv.it
- (2) Anglia Catholic Protection Services Ltd, Meadow View Industrial Estate, Reach Road, Burwell, Cambridgeshire, CB25 0GH, U.K.
- (3) Istituto Nazionale di Geofisica e Vulcanologia, via di Vigna Murata 605, 00143 Roma, Italy
- (4) Dipartimento di Scienze e Tecnologie, Università degli Studi di Camerino, Via Gentile III da Varano, 62032 Camerino, Italy

Abstract: The Piano di Pezza fault (PPF) is the north-westernmost segment of the >20 km long Ovindoli-Pezza active normal fault-system (central Italy). Although existing paleoseismic data document high vertical Holocene slip rates (~1 mm/yr) and a remarkable seismogenic potential of this fault, its subsurface setting and Pleistocene cumulative displacement are still unknown. We investigated the shallow subsurface of a key section of the PPF using seismic and electrical resistivity tomography coupled with time-domain electromagnetic measurements (TDEM). We provide 2-D Vp and resistivity images showing details of the fault structure and the geometry of the shallow basin infill down to 35-40 m depth. We can estimate the dip and the Holocene vertical displacement of the master fault. TDEM measurements in the fault hangingwall indicate that the pre-Quaternary carbonate basement may be found at ~90-100 m depth.

Key words: Central Apennines, refraction tomography, electrical resistivity tomography, time-domain electromagnetic measurement, active fault.

INTRODUCTION

The inner sector of the central Apennines of Italy is characterized by a complex network of Pliocene-Quaternary normal faults (Cavinato & de Celles, 1999; Ghisetti & Vezzani, 1999) that originated several intramontane continental basins (Bosi et al., 2003; Giaccio et al., 2012; Pucci et al., 2014). Under the current NE-directed extensional stress regime, the high crustal seismicity rate is mainly released by normal-faulting earthquakes (Chiarabba et al., 2005; Chiaraluca, 2012 and references). Paleoseismic analyses have unravelled the late Holocene slip-history of many active normal faults in this sector of the chain (Galli et al., 2008 and reference therein; Cinti et al., 2011). Notwithstanding the wealth of geological data constraining the recent evolution of most normal faults, only in few cases detailed subsurface information is available to provide a proper structural model of the basins they originated. Shallow geophysical investigation of active faults may extend the limited

investigation depth typical of paleoseismic trenching, moreover it can yield valuable information on long-term fault displacement and unravel minor structures (e.g.: Villani et al., 2014).

As a case-study, we selected the Piano di Pezza fault (hereinafter PPF, Figure 1). The PPF is the north-westernmost segment of the >20 km long active Ovindoli-Pezza normal fault system, and according to paleoseismic data it exhibits one of the highest Holocene vertical slip rates in the Apennines (~1 mm/yr; Pantosti et al., 1996), with large inferred vertical displacement per event (> 2 m). This fault bounds to the north a 5 km long continental basin (Piano di Pezza; Figure 1) filled with glacial, lacustrine and fluvial deposits, and its footwall exposes Mesozoic-Tertiary marine limestones. The subsurface structure and infilling thickness of this basin are unknown, nor the shallow geometry of the bounding fault is well constrained.



Figure 1: geological sketch of the Piano di Pezza basin. The blue lines indicate normal faults, the red lines thrusts.




Figure 2: close up view of the survey area showing seismic, electrical resistivity profiles and TDEM measure points (orange); the PPF exhibits two splays: we investigated the shallow subsurface of the lower one, where it displaces late Holocene alluvial fans; tA is trench A after Pantosti et al., 1996.



With the aim of extending information from previous paleoseismic analyses and very shallow GPR investigations (Jewell & Bristow, 2006), we performed an integrated geophysical survey focusing a key section of the PPF (Figure 2). We acquired 2 high-resolution seismic profiles, 1 electrical resistivity profile and 7 time-domain electromagnetic soundings (TDEM) crossing the fault where it displaces some small late Holocene alluvial fans, originating a ~5 m high fault scarp.

We try to infer the subsurface fault geometry and recover the depth of the pre-Quaternary basement.

METHODS AND DATA ACQUISITION

Traveltime tomography proved successful in imaging complex media, such as fault zones (Morey & Schuster, 1999; Sheley et al., 2003), so we conceived our seismic investigations to get tomographic images of the Piano di Pezza fault at depth. The 2 seismic profiles (named P1 and P2) are 142 m long each, and they were acquired using a 72 vertical geophones array (40 Hz eigenfrequency) with 2 m spacing and a sledgehammer as energy source. We handpicked ~20,000 first-arrival traveltimes that were input to a non-linear tomographic code to produce 2-D Vp models of the shallow subsurface (details of the method in: Improta et al., 2002).

Electrical resistivity tomography (ERT) is a powerful tool to investigate shallow faults and complex structures as well (Griffits & Barker, 1993; Storz et al., 2000). We acquired 1 ERT profile paralleling seismic line P2. We deployed a 64 electrodes and 315 m long array with both Wenner-Schlumberger and dipole-dipole configurations of quadrupoles. The apparent resistivity data were inverted through a linearized least-squares algorithm to obtain true resistivity sections (Constable et al. 1987; Loke & Dahlin 2002).




Figure 3: shallow electrical resistivity model across the Piano di Pezza fault.

In order to infer the pre-Quaternary basement depth in the hangingwall of the PPF, we also performed 5 time domain electromagnetic measurements (TDEM). The method is based on the induction of a time-varying secondary magnetic field produced by decay current in the ground. This secondary magnetic field is measured by an induction receiver coil generally placed inside of a transmitter loop. As the current diffuses deeper into the ground (at later time), the signal measurements provide information on the conductivity of the lower layers. Our data were collected using a high frequency 1D receiver

coil, placed at the centre and outside a transmitter square loop of 50 m size. We acquired very early time gates data starting from ~9 μ s (after current turn off) up to 800 μ s using a Geonics 47 transmitter. Random electromagnetic noise were also collected and checked against measured voltage data in order to discharge biased late time gates before data inversion. Data from field measurements were processed and inverted using SiTYEM\SEMDI software in order to obtain 1D resistivity models of the subsoil (Hydrogeophysics-Group, 2001)

PRELIMINARY RESULTS AND DISCUSSION

We show some preliminary results of our integrated geophysical surveys. The ERT model (Figure 3) with dipole-dipole configuration across the fault scarp clearly depicts the ~50° SW-dipping fault zone involving alluvial fan deposits with resistivity <500 Ω m and their concurrent thickening in the hangingwall. Moreover, the top-surface of the underlying high-resistivity body (>1000 Ω m) shows an evident dip towards the fault, suggesting back-tilting due to fault activity. We interpret the shallower layers as Holocene alluvial fan gravelly and sandy deposits (recovered in the trenches by Pantosti et al., 1996) laying over a > 20 thick Last Glacial Maximum till (~21-18 ka; Cassoli et al., 1986; Giraudi, 1989). Vp tomograms suggest this alluvial fan is affected by > 10 m vertical displacement (Figure 4). They also help defining the fault-zone down to 35-40 m depth suggesting it has a ~50° dip.




Figure 4: detail of the Vp tomogram of seismic line P2 across the PPF compared with trench A (after Pantosti et al., 1996).

TDEM data inversion yielded 1-D vertical resistivity models (Fig. 5): they indicate that ~200 m far from the surface fault trace the pre-Quaternary carbonate basement is 90-100 m deep, thus giving a robust constraint to the evaluation of the cumulative offset of the PPF (~150 m). Moreover, for the uppermost part of the subsoil, TDEM and ERT model significantly agree in terms of depth and thicknesses of the shallower resistivity layers.




Figure 5: 1-D resistivity models obtained through inversion of TDEM data (position of soundings along the ERT profile is indicated by red stars). A deep resistive layer ($>5000 \Omega\text{m}$), interpreted as the carbonate basement, is evidenced at 90-100 m depth (beneath soundings 2offB, 2c and 2offA) in the hangingwall of the PPF. Data from soundings 1off and 1c are less reliable in the deeper part due to strong 3D effects.

Our geophysical data yield for the first time valuable information on the subsurface setting of an important active normal fault exhibiting one of the highest Holocene slip rates in the central Apennines. These data are crucial to understand the middle- to long-term behaviour of this fault, its overall displacement and possibly its inception age.

Acknowledgements: This study was performed in the framework of the FIRB Abruzzo project, "High-resolution analyses for assessing the seismic hazard and risk of the areas affected by the 6 April 2009 earthquake" (<http://progettoabruzzo.rm.ingv.it/en>; P. Villani, V. Sapia & R. Civico funding code: RBAP10ZC8K_005; F. Baccheschi funding code: RBAP10ZC8K_003; G. Di Giulio & M. Vassallo funding code: RBAP10ZC8K_007).

References

- Bredosian, P.A., M.K. Burgess & T. Nishikawa, (2013). Faulting and groundwater in a desert environment: constraining hydrogeology using time-domain electromagnetic data. *Near Surface Geophysics*, 11, (5), 545-555.
- Bosi, C., F. Galadini, B. Giaccio, P. Messina, & A. Sposato, (2003). Plio-Quaternary continental deposits in the Latium-Abruzzi Apennines: the correlation of geological events across different intermontane basins. *Il Quaternario*, 16 (1Bis), 55-76.
- Cassoli, A., L. Corda, C. Lodoli, A. Malatesta, M.V. Molaroni, & A. Ruggeri, (1986). Il glacialismo quaternario del gruppo Velino-Ocre-Sirente. *Memorie Società Geologica Italiana*, 35, 855-867.
- Cavinato, G.P., & P.G. De Celles, (1999). Extensional basins in the tectonically bimodal central Apennines fold-thrust belt, Italy: response to corner flow above a subducting slab in retrograde motion. *Geology*, 27, 955-958.
- Chiarabba, C., L. Jovane & R. Di Stefano, (2005). A new view of Italian seismicity using 20 years of instrumental recordings. *Tectonophysics*, 305, 251-268.
- Chiaraluce, L., (2012). Unraveling the complexity of Apenninic extensional fault systems: A review of the 2009 L'Aquila earthquake (Central Apennines, Italy). *Journal of Structural Geology*, 42, 2-18.
- Cinti, F.R., D. Pantosti, P.M. De Martini, S. Pucci, R. Civico, S. Pierdominici, L. Cucci, C.A. Brunori, S. Pinzi, & A. Patera, (2011). Evidence for surface faulting events along the Paganica fault prior to the 6 April 2009 L'Aquila earthquake (central Italy). *Journal of Geophysical Research*, 116, 2156-2202.
- Constable, S.C., R.L. Parker & C.G. Constable, (1987). Occam's inversion: A practical algorithm for generating smooth models from electromagnetic sounding data. *Geophysics*, 52, (3), 289-300.
- Galli, P., F. Galadini & D. Pantosti, (2008). Twenty years of paleoseismology in Italy: *Earth Sciences Review*, 88, 89-117.
- Giaccio, B., P. Galli, P. Messina, E. Peronace, G. Scardia, G. Sottili, A. Sposato, E. Chiarini, B. Jicha & S. Silvestri, (2012). Fault and basin depocentre migration over the last 2 Ma in the L'Aquila 2009 earthquake region, central Italian Apennines. *Quaternary Science Review*, 56, 69-88.
- Ghissetti, F. & L. Vezzani, (1999). Depths and modes of Pliocene-Pleistocene crustal extension of the Apennines (Italy). *Terra Nova*, 11, 67-72.
- Giaccio, B., P. Galli, P. Messina, E. Peronace, G. Scardia, G. Sottili, A. Sposato, E. Chiarini, B. Jicha & S. Silvestri, (2012). Fault and basin depocentre migration over the last 2 Ma in the L'Aquila 2009 earthquake region, central Italian Apennines. *Quaternary Sci. Rev.* 56, 69-88, doi: 10.1016/j.quascirev.2012.08.016.
- Giraudi, C., (1989). Datazione con metodi geologici delle scarpate di faglia post-glaciali di Ovindoli - Piano di Pezza (Abruzzo - Italia Centrale): implicazioni. *Memorie Società Geologica Italiana*, 42, 29-39.
- Griffiths, D.H. & R.D. Barker, (1993). Two-dimensional resistivity imaging and modelling in areas of complex geology. *J. Appl. Geophys.* 29, 211-226.
- Hydrogeophysics Group, (2001). Getting Started with SiTEM and SEMDI. University of Aarhus. Aarhus, Denmark.
- Improta, L., A. Zollo, A. Herrero, R. Frattini, J. Virieux, & P. Dell'Aversana, (2002). Seismic imaging of complex structures by non-linear travelttime inversion of dense wide-angle data: Application to a thrust belt. *Geophysical Journal International*, 151, 264-278.
- Jewell, C.J., & C. Bristow, (2006). GPR studies in the Piano di Pezza area of the Ovindoli - Pezza fault, central Apennines, Italy: extending paleoseismic trench investigations with high-resolution GPR profiling. *Near Surface Geophysics*, 4, (3), 147-153.
- Loke, M.H. & T. Dahlin, (2002). A comparison of the Gauss-Newton and quasi-Newton methods in resistivity imaging inversion. *Journal of Applied Geophysics*, 49, 149-162.
- Morey, D. & G.T. Schuster, (1999). Paleoseismicity of the Oquirrh fault, Utah, from shallow seismic tomography. *Geophysical Journal International*, 138, 25-35, doi: 10.1046/j.1365-246x.1999.00814.x.
- Pantosti, D., G. D'Addezio & F.R. Cinti, (1996). Paleoseismicity of the Ovindoli-Pezza fault, central Apennines, Italy: a history including a large, previously unrecorded earthquake in the Middle Ages (860-1300 A.D.). *Journal of Geophysical Research*, 101, 5937-5959.
- Pucci, S., F. Villani, R. Civico, D. Pantosti, P. Del Carlo, A. Smedile, P.M. De Martini, E. Pons-Branchu & A. Gueli, (2014). Quaternary geology map of the Middle Aterno Valley, 2009 L'Aquila earthquake area (Abruzzi Apennines, Italy). *Journal of Maps*. doi: 10.1080/17445647.2014.927128.
- Sheley, D., T. Crosby, M. Zhou, J. Giacomina, J. Yu, R. He, & G.T. Schuster, (2003). 2-D seismic trenching of colluvial wedges and faults. *Tectonophysics*, 368, 51-69.
- Storz, H., W. Storz & F. Jacobs, (2000). Electrical resistivity tomography to investigate geological structures of the Earth's upper crust. *Geophysical Prospecting*, 48, 455-471. doi: 10.1046/j.1365-2478.2000.00196.x.
- Villani, F., S. Pucci, R. Civico, P.M. De Martini, I. Nicolosi, F. D'AJello Caracciolo, R. Carluccio, G. Di Giulio, M. Vassallo, A. Smedile & D. Pantosti, (2014). Imaging the structural style of an active normal fault through multi-disciplinary geophysical investigation: a case study from the Mw 6.1, 2009 L'Aquila earthquake region (central Italy). *Geophysical Journal International*, 200 (3), 1676-1691. doi: 10.1093/gji/ggu462.